



Alstorisine A, a *nor*-monoterpenoid indole alkaloid from cecidogenous leaves of *Alstonia scholaris*



Ying-Ying Chen^{a,b,c}, Jing Yang^b, Xing-Wei Yang^b, Afsar Khan^{b,d}, Lu Liu^{b,c}, Bei Wang^{b,c}, Yun-Li Zhao^b, Ya-Ping Liu^b, Zhong-Tao Ding^{a,*}, Xiao-Dong Luo^{a,b,*}

^a Ministry of Education, School of Chemical Science and Technology, Yunnan University, Kunming 650091, People's Republic of China

^b State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Science, Kunming 650201, People's Republic of China

^c University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China

^d Department of Chemistry, COMSATS Institute of Information Technology, Abbottabad 22060, Pakistan

ARTICLE INFO

Article history:

Received 11 December 2015

Revised 3 March 2016

Accepted 8 March 2016

Available online 8 March 2016

Keywords:

Alstonia scholaris

Galls

Monoterpenoid indole alkaloid

Absolute configuration

ABSTRACT

Alstorisine A (**1**), a novel *nor*-monoterpenoid indole alkaloid with 6/5/6/6 fused-ring system, regarded as a key intermediate from melodinine E to mersicarpine, was isolated from the cecidogenous leaves of *Alstonia scholaris*. Its structure was identified on the basis of extensive spectroscopic analysis and the comparison of experimental and calculated ECDs. A plausible biogenesis was also postulated.

© 2016 Elsevier Ltd. All rights reserved.

Plants defend against insect herbivory for their survival, while some insects spawn on the leaves instead of feeding them immediately so that their eggs can survive by sojourn in these leaves. Then, some special tissues more like tumor came into being, which were called galls and were induced by gall-formers such as insects and fungi on host plants. The chemical constituents of the cecidogenous plants may have some difference with the normal ones.¹ During the years of observation on *Alstonia scholaris*, we found that the leaves of *A. scholaris* may suffer galls on surface.² We wondered whether the constituents would have some changes due to the galls on the leaves. This inspired us to compare the difference between normal leaves and cecidogenous leaves. The HPLC fingerprint analysis of the extracts of two forms showed some obvious difference, that is, the contents of the main components changed and some trace compositions were produced (see Fig. 1 and Supplementary data). More importantly, cecidogenous leaves of *A. scholaris* seem not to have been studied chemically before. Motivated by the interesting phenomenon, intriguing architectures,^{3,7} and bioactivities⁴ of monoterpenoid indole alkaloids in this plant, a phytochemical study of the cecidogenous leaves was carried out. As a result, a *nor*-monoterpenoid indole alkaloid, alstorisine

A (**1**) with a 6/5/6/6 fused-ring system was isolated. Furthermore, it was not distributed in healthy leaves, but appeared equally both in galls and the normal part of cecidogenous leaves at a concentration of 3 µg/g, which was indicated by LC–MS quantitative analysis (see Supplementary data). In alstorisine A, a carbon (C-5) of the skeleton was degraded compared to melodinine E, scholarisine G, and leuconoxine, which have attracted much attention for their total synthesis and asymmetric cascade catalysis.⁵ Zhu and co-workers developed a unified strategy for the enantioselective synthesis of leuconolam-leuconoxine-mersicarpine subfamily of Aspidosperma alkaloids, while melodinine E served as springboard to reach leuconoxine, leuconodine A, leuconodine C, leuconodine F, and leuconolam.^{5g} Then, alstorisine A (**1**) could be regarded as a key intermediate from melodinine E to mersicarpine.^{5g} Besides, 20 known compounds were also isolated and identified as 4-aminobenzylamine, 3-(aminomethyl)-benzoic acid, (Z)-16-formyl-5 α -methoxystrictamine,⁶ 5 α -methoxystrictamine,⁷ picralinal,⁸ picrinine,⁸ (+)-geissoschizine,⁹ (+)-vincadifformine,¹⁰ strictamine,¹¹ vallesiachotamine,¹² isovallesiachotamine,¹² vallesamine,¹³ akuammidine,¹⁴ tubotaiwine,¹⁵ tubotaiwine *N*-oxide,¹⁶ burnamine,¹⁷ scholarine,¹⁸ (19R)-scholaricine,¹⁹ (19S)-scholaricine,¹⁹ and hedyotisol-A.²⁰ The new compound was tested for its bioactivity of regulating hippocampal neural stem cells (NSCs) proliferation in vitro. Reported herein is the isolation, structural elucidation, and proposed biogenic pathway of **1**.

* Corresponding authors. Tel.: +86 871 65223177.

E-mail addresses: ztding@ynu.edu.cn (Z.-T. Ding), xdluo@mail.kib.ac.cn (X.-D. Luo).




Figure 1. Fingerprint of the tested samples (black: total alkaloids from the cecidogenous leaves of *A. scholaris*; red: total alkaloids from the normal leaves of *A. scholaris*).

Air-dried and powdered leaves with galls of *A. scholaris* (10 kg) were extracted three times with MeOH under reflux conditions and the solvent was evaporated in vacuo. The extract was dissolved in 0.3% HCl, and the solution was filtrated and subsequently basified to pH 9–10, using ammonia. The solution was partitioned with EtOAc, affording aqueous and EtOAc phases. The EtOAc fraction (113 g) was chromatographed on a silica gel column, eluting with CHCl₃–MeOH [from 30:1 to 0:1], to afford seven fractions (I–VII). Fraction II (18.5 g) was subjected to a preparative reversed phase C₁₈-MPLC column with a gradient flow of 30–80% (v/v) aqueous MeOH to yield six subfractions II-1–II-6. Subfraction II-5 (2.1 g) was further separated by silica gel to get the subfraction containing the target molecule. Compound **1** (20 mg) was purified from this subfraction via Prep-TLC using CHCl₃–MeOH (15:1) as an eluent.

Compound **1**²¹ (Fig. 2) was isolated as a white amorphous powder and gave a positive reaction with Dragendorff's reagent. Its molecular formula was obtained as C₁₈H₂₂N₂O₂ by HRESIMS (*m/z* 299.1756 [M+H]⁺) in association with ¹H and ¹³C NMR data, which indicated 9 degrees of unsaturation. Its IR spectrum showed characteristic absorption bands at 3432 cm⁻¹ for hydroxyl group, 1631 cm⁻¹ for lactam group, and 1474 and 1099 cm⁻¹ for an aromatic ring. The UV spectrum showed absorption maxima characteristic of β-anilinoacrylate chromophore (337, 264, 245, and 210 nm).²² The ¹H NMR spectrum revealed the existence of a 1,2,4-trisubstituted benzene ring [δ_{H} 6.92 (1H, d, *J* = 2.3 Hz, H-9), 6.80 (1H, dd, *J* = 8.7, 2.3 Hz, H-11), 8.14 (1H, d, *J* = 8.7 Hz, H-12)]. The correlations of δ_{H} 8.14 (d, *J* = 8.7 Hz, H-12) with 6.80 (dd, *J* = 8.7, 2.3 Hz, H-11) in the ¹H–¹H COSY spectrum, as well as the

correlations of δ_{H} 6.92 (d, *J* = 2.3 Hz, H-9) with δ_{C} 142.2 (C-7) in the HMBC spectrum, suggested that C-10 (δ_{C} 153.6) was substituted by a –OH function. The ¹³C NMR (BB and DEPT) spectra of **1** displayed a total of 18 carbon resonances which were assigned to one methyl (δ_{C} 7.2), seven methylenes (δ_{C} 105.8, 39.3, 30.7, 29.3, 27.1, 23.9, and 20.2), three methines (δ_{C} 119.2, 116.9, and 107.1), and seven quaternary carbons (δ_{C} 169.2, 153.6, 142.2, 135.5, 128.8, 84.8, and 37.2) (Table 1). The spectral data as well as a series of monoterpene indole alkaloids (MIAs) isolated from *A. scholaris* suggested that **1** might be a MIA derivative. Unlike the other intact MIAs, in the HMBC spectrum of compound **1**, the correlations from δ_{H} 5.69 (s, H_a-6) and 5.29 (s, H_b-6) to δ_{C} 128.8 (C-8),

Table 1
¹H (600 MHz), ¹³C (150 MHz) NMR, and HMBC data of **1** (δ in ppm) in CDCl₃

No.	¹ H, mult. (<i>J</i> Hz)	¹³ C, type	HMBC
2		169.2, s	
3	3.11, td (12.1, 3.3) 2.75, overlap	39.3, t	15, 21
6	5.69, s 5.29, s	105.8, t	7, 8, 21
7		142.2, s	
8		128.8, s	
9	6.92, d (2.3)	107.1, d	7, 10, 11, 13
10		153.6, s	
11	6.80, dd (8.7, 2.3)	116.9, d	9, 10, 13
12	8.14 d (8.7)	119.2, d	8, 10, 13
13		135.5, s	
14	1.87, overlap 1.55, m	20.2, t	20
15	1.86, overlap 1.74, m	30.7, t	3, 14, 17, 19, 20, 21
16	2.74, overlap 2.46, m	29.3, t	2, 17, 20
17	2.75, overlap 1.48, m	23.9, t	2, 15, 19, 20, 21
18	0.79, t (7.5)	7.2, q	19, 20
19	1.44, q (7.5) 1.37, q (7.5)	27.1, t	15, 17, 18, 20, 21
20		37.2, s	
21		84.8, s	




Figure 2. Structure of alstorisine A (**1**).

Download English Version:

<https://daneshyari.com/en/article/5259241>

Download Persian Version:

<https://daneshyari.com/article/5259241>

[Daneshyari.com](https://daneshyari.com)