



Actinide (Th⁴⁺ and UO₂²⁺) assisted oxidative coupling of *ortho*-phenylenediamine in the presence of oxygen



Branson A. Maynard, Charmaine D. Tutson, K. Sabrina Lynn, Chasity W. Pugh, Anne E. V. Gorden*

Auburn University, 179 Chemistry Building, Auburn, AL 36849, USA

ARTICLE INFO

Article history:

Received 30 October 2015

Revised 7 December 2015

Accepted 12 December 2015

Available online 14 December 2015

Keywords:

ortho-Phenylenediamine

Catalysis

Thorium

Actinide

Diaminophenazine

ABSTRACT

While the actinide salts because of their Lewis acidity and flexible coordination geometry demonstrate unique possibilities as catalysts, research with actinide catalysts typically involves complexed organoactinide frameworks. Many actinide salts have yet to be explored in catalysis or as reagents in metal mediated synthesis. Here, the synthesis of the heterocycle 2,3-diaminophenazine has been reported in good yields in the presence of oxygen via the first use of thorium nitrate or uranyl nitrate as the catalyst.

© 2015 Elsevier Ltd. All rights reserved.


Introduction

Recently, the early 5f elements have been of interest for their potential use in catalysis, metal mediation, and stoichiometric reactivity.^{1–6} Actinides have 5f valence orbitals and large ionic radii, both of which offer distinguishing chemical characteristics that can be exploited in catalysis.⁷ Reports detailing some of these synthetic methods^{3,4,8,9} or catalytic reactions^{2,5,10,11} incorporating actinide metals have become more common as new starting materials have been developed.^{12–14} For example, thorium catalysts have been used for the cyclization and hydroalkoxylation of alkynyl alcohols² and thorium and uranium Cp* complexes have been used in the ring-opening metathesis polymerization of cyclic esters.¹¹ The focus of this work is typically based on an actinide center saddled with sterically hindering ligands (such as cyclopentadiene,¹³ pentamethylcyclopentadienyl, or *N*-ethylmethylamine^{11,15}), although U_I₃ has been reported to catalyze Diels–Alder and Mukaiyama aldol reactions.^{16,17} Ultimately, the ligand and the metal are bound covalently to some degree, and the reactivity of these ligand–actinide complexes and their mechanisms of action have not been fully explained.^{11,15} For example, Arnold and coworkers have used an organothorium metallacycle precatalyst, [(Me₃Si)₂N]Th[k₂-(N,C)-CH₂Si(CH₃)N(SiMe₃)], to regioselectively oligomerize terminal alkynes producing organic enynes and this demonstrates the first example of using a Th catalyst such as this in C–C bond formation.¹⁸ Although all of these methods have aided

in the understanding of the chemistry of the actinides, they require laborious preparation and, being either air or moisture sensitive, have short shelf-lives. In contrast to these organoactinide catalysts, actinide salts offer longer shelf-lives and increased ease of use. They are commercially available, and still remain to be characterized.

The heterocycle 2,3-diaminophenazine (DAP), is of interest in the self-assembly of nanobelts,¹⁹ and it is typically prepared via H₂O₂ catalysis with horseradish peroxidase and fungal laccase enzymes.²⁰ Li took advantage of the electroactivity of DAP when using the rapid oxidation of OPD to DAP via cupric ion-catalysis to synthesize electroactive nanoparticles that were to dope a graphene nanolabel. This graphene nanolabel was then coupled with a peptide-based probe to target SMO, a protein connected to metastatic tumor cells that is used as a biomarker, to determine the metastatic activity of tumor cells.^{21,22} Non-biological synthetic pathways are known, including catalysis by Fe(III), Co(II), and Cu(II) metal salts.^{19,23–25} We noted that the unencumbered 5f orbitals of actinide salts may offer another non-biological pathway to catalyze this synthesis. Here, we describe the thorium nitrate, uranyl nitrate, and oxygen mediated synthesis of 2,3-diaminophenazine from *ortho*-phenylenediamine (OPD) (see Scheme 1). Of note, simple, sterically unencumbered actinide centers have rarely been reported in the literature.^{17,18} Few reactions have been reported with simple actinide salts,^{17,18,26} and no catalytic reactions have been reported using simple thorium salts.

* Corresponding author.



Scheme 1. Oxidation of *ortho*-phenylenediamine to 2,3-diaminophenazine.




Figure 1. Percent yield of oxidative coupling of *ortho*-phenylenediamine versus time.




Figure 2. Percent yield of oxidative *ortho*-phenylenediamine coupling with varying conditions.

Results and discussion

As shown in Figure 1, dioxygen alone can oxidize *ortho*-phenylenediamine to 2,3-diaminophenazine. Incorporating the simple actinide nitrate salts [$\text{Th}(\text{NO}_3)_4$ or $\text{UO}_2(\text{NO}_3)_2$] as a catalyst with just 1% catalyst loading, enhances the oxidation of *ortho*-phenylenediamine to 2,3-diaminophenazine, by factors of 2 and 1.5, respectively, in the presence of dioxygen. While this oxidation occurs much slower than results found using simple FeCl_3 salt—in that case 80% catalyst loading was required (0.48 mmol FeCl_3 to 0.60 mmol of *ortho*-phenylenediamine).¹⁹ Here, only a 1% catalyst loading with $\text{Th}(\text{NO}_3)_4$ or $\text{UO}_2(\text{NO}_3)_2$ is required. It is evident that the addition of actinide ions (Th^{4+} and UO_2^{2+}) to the aqueous solution provides metal mediation for the oxidative generation of 2,3-diaminophenazine. Increasing the Th^{4+} concentration by a factor of 2 increases the formation of the 2,3-diaminophenazine product as seen in Figure 1 (Th and Th run 2). This indicates that the inclusion the actinide (Th^{4+} or UO_2^{2+}) ions in the presence of dioxygen exhibits a synergistic behavior, as the sum of the ‘ O_2 ’ and degassed

actinide salt reactions is less than the isolated product in the ($\text{Th}^{4+} + \text{O}_2$) reaction alone.

To better understand the influence of dioxygen in the mechanism of the oxidation reaction, two reactions were prepared and allowed to react for seven days. In the first, a Schlenk flask containing water and *ortho*-phenylenediamine with $\text{Th}(\text{NO}_3)_4$ as the actinide source was degassed by bubbling Argon using a freeze–pump–thaw method to remove as much dioxygen as possible. In the second reaction, a Schlenk flask containing water and *ortho*-phenylenediamine with $\text{UO}_2(\text{NO}_3)_2$ as the actinide source was also degassed using freeze–pump–thaw. After 7 days, no observable precipitate was present in the reaction flask containing $\text{Th}(\text{NO}_3)_4$, and no color change was observed. No phenazine was detected upon workup and analysis. The reaction flask with $\text{UO}_2(\text{NO}_3)_2$ was found to contain 5.2% yield of isolable 2,3-diaminophenazine. This result indicates that the presence of dioxygen, is needed for the oxidation to occur effectively.¹⁶

A third reaction scheme was initiated to further determine how vital oxygen was to the reaction scheme. A Schlenk flask was set up with only a stir bar in an aqueous solution of *ortho*-phenylenediamine. Argon was bubbled through the solution for two hours, and then put through three cycles of freeze–pump–thaw in an effort to remove all possible traces of O_2 . This resulted in a 25% yield of 2,3-diaminophenazine. Thus, it is evident that an oxygen donor present in the reaction mixture participates in the reaction pathway. This result would require the oxygen from H_2O to serve as the oxidant. Interestingly, as this reaction contained only H_2O and *ortho*-phenylenediamine, a literature search was unable to find any previous Letters on the temperature induced oxidation of *ortho*-phenylenediamine in an inert atmosphere.

The effect of light on this reaction was explored by running the highest yielding actinide salt reaction, with $\text{Th}(\text{NO}_3)_4$, in the absence of light. The reaction was set up in a 250 mL round bottom flask and heated at 80 °C for 6 days while wrapped in aluminum foil to block out light. As shown in Figure 2, the absence of light did not completely quench the reaction and a 33% yield was obtained as compared to the reaction in the presence of light in which a yield 92% product was obtained. Thus, as with the actinide and the O_2 previously discussed, light is critical to optimizing the reaction, indicating that there are likely multiple concurrent reaction pathways and one of these is likely through a radical pathway.

To determine whether or not the reaction proceeds through interactions with the pi orbitals of the *ortho*-phenylenediamine or through interactions with lone pairs of the amines present on the *ortho*-phenylenediamine, various substrates were subjected to the same parameters as the highest yielding actinide catalyzed reaction. Table 1 outlines the substrates that were explored to determine if the $\text{Th}(\text{NO}_3)_4$ would also oxidatively couple them under these same conditions. To test the effect of the pi orbitals, reactions with cyclohexene, benzene, and 1,4-cyclohexadiene were examined. To determine the effect of the lone pairs, aniline, and catechol were used. Substrates were also reacted in the presence of *ortho*-phenylenediamine to determine if the oxidative coupling of *ortho*-phenylenediamine was favored. In the presence of *ortho*-phenylenediamine, the only product found during analysis was 2,3-diaminophenazine, in yields up to 64%. Thorium(IV) nitrate was proven futile as a catalyst to oxidatively couple the substrates with themselves. Mass spec analysis supported that the reaction between *ortho*-phenylenediamine with aniline and *ortho*-phenylenediamine with catechol were catalyzed. In both cases, however, 2,3-diaminophenazine was also synthesized and NMR analysis did not support the presence of the aniline/*ortho*-phenylenediamine product. Thus, for thorium(IV) nitrate reactions, what was observed is better suited to electron rich environments.

Download English Version:

<https://daneshyari.com/en/article/5261401>

Download Persian Version:

<https://daneshyari.com/article/5261401>

[Daneshyari.com](https://daneshyari.com)