

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

Tetrahedron Letters

journal homepage: www.elsevier.com/locate/tetlet

Displacement-based, chromogenic calix[4]pyrrole–indicator complex for selective sensing of pyrophosphate anion

Sandeep Kaur^a, Hoon Hwang^a, Jeong Tae Lee^{b,*}, Chang-Hee Lee^{a,*}^a Department of Chemistry, Kangwon National University, Chun Cheon 200-701, Republic of Korea^b Department of Chemistry, Hallym University, Chun Cheon 200-700, Republic of Korea

ARTICLE INFO

Article history:

Received 2 April 2013

Revised 23 April 2013

Accepted 30 April 2013

Available online xxxx

Keywords:

Calix[4]pyrrole

Indicator displacement assay

Pyrophosphate

Azophenolate

Anion recognition

ABSTRACT

A supramolecular complex composed of bis-pyridinium picket calix[4]pyrrole and azophenol indicator is a highly visible colorimetric displacement assay and sensor. The system shows significant selectivity and a higher affinity for pyrophosphate anions over other competing anions.

© 2013 Elsevier Ltd. All rights reserved.


The design and synthesis of sensitive chemosensors for selective detection of anions¹ have received considerable attention in the chemical, biological, and environmental sciences.² The key components in an effective chemosensor are the recognition domain (binding site) and the signaling unit (indicator).³ In most molecularly constructed sensors, the receptor site and signaling unit are covalently linked to facilitate observable changes associated with the binding interaction. Anslyn and co-workers⁴ have developed the indicator-displacement assay (IDA), where sensing of a target analyte is achieved by a binding-induced displacement within a supramolecular receptor–indicator complex. This approach relies upon the competition between the analyte and the indicator for a binding cavity on the receptor; the analyte must have the higher binding affinity. It has been used for the selective sensing of anions such as phosphate,⁵ pyrophosphate,⁶ nitrate,⁷ cyanide,⁸ and citrate.⁹

The detection of pyrophosphate anion ($\text{HP}_2\text{O}_7^{3-}$) is particularly important for the analysis of bioenergetic and metabolic processes.¹⁰ It plays a critical role in energy storage¹¹ and signal transductions, as well as being a structural component of teeth and bones. Also, it is a product of ATP hydrolysis and participates in many enzymatic reactions¹² such as the adenylate cyclase-catalyzed synthesis of cyclic AMP, aminoacyl tRNA synthetase-catalyzed attachment of amino acids to tRNA in protein synthesis, and DNA sequencing/replication¹³ catalyzed by DNA polymerase. High levels of $\text{HP}_2\text{O}_7^{3-}$ are known to cause several diseases.¹⁴

Zinc–dipicolylamine (Zn(II)–DPA) and Cu(II)–DPA complexes have been employed as IDAs for the detection of pyrophosphate anions.^{15,16} The metal centers become coordination spheres to accommodate the oxoanions. Other receptors for pyrophosphate anion include macrocyclic pyrrole, imidazolium-based macrocycles, and dipyrrolyquinoxalines.¹⁷ We reported a bis-pyridinium calix[4]pyrrole derivative for ‘turn on’ fluorescence detection of pyrophosphate in an aqueous organic solvent¹⁸ that utilizes a hydrogen bonding interaction and electrostatic interactions combined with a fluorescent dye-displacement assay. We have also developed a supramolecular receptor–indicator complex¹⁹ composed of bis-pyridinium calix[4]pyrrole and an azo dye for selective recognition of $\text{HP}_2\text{O}_7^{3-}$ over other competing anions, including F^- and AcO^- . The recognition of F^- in organic media was achieved with a colorimetric IDA using an octamethyl-calix[4]pyrrole-(*p*-nitrophenolate) complex by Sessler and co-workers²⁰ and a merocyanine dye by Machado and co-workers.²¹

Here, we report on an IDA-based colorimetric detection of $\text{HP}_2\text{O}_7^{3-}$ anion using dicationic calix[4]pyrrole combined with an azo dye indicator. The pyrrole was designed to allow multiple interactions with the guest anion (hydrogen bonding, anion– π interactions, and coulombic interactions). The azo dye, initially bound to the receptor, is replaced by the target analyte, resulting in colorimetric detection (Scheme 1). The *cis*-5,15-(4-pyridyl)-5,10,10,15,20,20-hexamethylcalix[4]pyrrole was prepared in moderate yield by acid-catalyzed condensation of 5-(4-pyridyl)dipyrromethane with acetone. The hexafluorophosphate salt of bis-pyridinium calix[4]pyrrole **1** was obtained via methylation of *cis*-5,15-(4-pyridyl)-5,10,10,15,20,20-hexamethylcalix[4]pyrrole,

* Corresponding authors. Tel.: +82 33 250 8490; fax: +82 33 253 7582 (C.-H.L.).
E-mail address: chhlee@kangwon.ac.kr (C.-H. Lee).



Scheme 1. Formation of the complex $[1\cdot 2^-]$ and the recognition of $\text{HP}_2\text{O}_7^{3-}$ via displacement of dye 2^- .

followed by treatment with NH_4PF_6 . Tetrabutylammonium salt of azo-dye 2^- was obtained in good yield by treating 4-(4-nitrophenylazo)resorcinol with tetrabutylammonium hydroxide. The structure of all the compounds was confirmed with spectroscopic means (SI).

The anionic form of indicator 2^- ($10\ \mu\text{M}$) has a red/pink color with twin absorption bands at $538\ \text{nm}$ ($\epsilon_{\text{max}} = 42,800\ \text{l}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$) and $447\ \text{nm}$ ($\epsilon_{\text{max}} = 49,500\ \text{l}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$) in acetonitrile. Incremental addition of receptor 1 (0 – $30\ \mu\text{M}$) to the solution of 2^- ($10\ \mu\text{M}$) results in decreased absorbance at $538\ \text{nm}$ and increased absorbance at $447\ \text{nm}$, with a $10\ \text{nm}$ bathochromic shift, as shown in Figure 1.

The binding of indicator 2^- to receptor 1 is accompanied by a distinctive color change from pink to yellow. Saturation occurs when $30\ \mu\text{M}$ of receptor is added (Fig. 2). The isosbestic point at $500\ \text{nm}$ indicates an equilibrium complexation of receptor 1 and indicator 2^- . A Job plot also supports the 1:1 binding stoichiometry between receptor 1 and indicator 2^- .²² Analysis of the titration data in Figure 1 with a non-linear least square fit (HypSpec²³) yields an association constant $K_a = (1.80 \pm 0.04) \times 10^6\ \text{M}^{-1}$. These observations clearly indicate that formation of the supramolecular receptor–indicator complex $[1\cdot 2^-]$ is favorable and that the hydrogen-bonding interactions between the anionic indicator 2^- and the pyrrole N–H bonds are strong. The binding behavior of the anionic indicator 2^- and receptor 1 can be monitored by ^1H NMR spectroscopy (Fig. 3). When receptor 1 ($2.28\ \text{mM}$) is titrated with anionic




Figure 2. Job plot for the binding between 1 and 2^- in CH_3CN .

indicator 2^- in CD_3CN , the signals corresponding to the *ortho*-protons from the anionic center are shifted up-field and the signals from the rest of the aromatic protons become broad relative to the spectrum of pure 2^- in CD_3CN .

In addition, the signal corresponding to NH protons of receptor 1 shows a considerable down-field shift from $\delta\ 7.90$ to $\delta\ 10.85\ \text{ppm}$ (Fig. 3 (iii)), indicating hydrogen bonding between phenolate anion and pyrrole N–H bonds, as well as possible π – π interactions.²⁴ Only 1.0 equiv of 2^- is required for complete binding.

The anion recognition properties of the complex $[1\cdot 2^-]$ were investigated by UV–vis absorption spectroscopy in CH_3CN . When it is titrated with the anions F^- , CN^- , AcO^- , and Cl^- (as their tetrabutyl ammonium salt, 0 – $17.5\ \mu\text{M}$), only small increases in absorbance at $538\ \text{nm}$ are observed (Fig. 4). However, a significant change in absorbance, as well as in the visual color, is observed upon titration with pyrophosphate anion ($\text{HP}_2\text{O}_7^{3-}$). Thus the affinity of pyrophosphate anion toward receptor 1 is strong and is capable of complete replacement of the indicator to form the new complex $[1\cdot\text{HP}_2\text{O}_7^{3-}]$. In contrast, H_2PO_4^- , Br^- , and I^- anions do not produce appreciable changes in the absorption spectra.

However, the initial $[1\cdot 2^-]$ absorption spectrum is shifted to a broad absorption band at 400 – $430\ \text{nm}$ upon titration with HSO_4^- ($17.5\ \mu\text{M}$). This spectral change is associated with the ionization of monobasic HSO_4^- to dibasic SO_4^{2-} by indicator anion 2^- , which is sufficiently basic to deprotonate HSO_4^- . Formation of the resulting azophenol derivative 2 can be confirmed by comparing the absorption spectra with 2^- . The detection limit²⁵ for $\text{HP}_2\text{O}_7^{3-}$ is




Figure 1. Changes in the absorption spectrum of $1\cdot 2^-$ ($10\ \mu\text{M}$) upon titration with pyrophosphate ion (0 – $30\ \mu\text{M}$) in CH_3CN . The inset displays the color of the $[1\cdot 2^-]$ and the complex $[1\cdot\text{HP}_2\text{O}_7^{3-}]$.

Download English Version:

<https://daneshyari.com/en/article/5264155>

Download Persian Version:

<https://daneshyari.com/article/5264155>

[Daneshyari.com](https://daneshyari.com)