



N-arylation of indoles with aryl halides using copper/glycerol as a mild and highly efficient recyclable catalytic system



Dilip Kumar T. Yadav, Sanil S. Rajak, Bhalthandra M. Bhanage*

Department of Chemistry, Institute of Chemical Technology, N. Parekh Marg, Matunga, Mumbai 400019, India

ARTICLE INFO

Article history:

Received 1 November 2013

Revised 11 December 2013

Accepted 14 December 2013

Available online 21 December 2013

Keywords:

Aryl halides

Arylation

NH-heterocycles

Copper

Glycerol

ABSTRACT

A simple, efficient, and inexpensive copper catalyzed protocol has been developed for cross-coupling of indoles with aryl halides by using glycerol as a green recyclable solvent and DMSO as an additive. The reaction was carried out at low catalyst loading furnishing various *N*-aryl indoles in good to excellent yield. This copper/glycerol mixture was recycled up to four runs without any loss in its catalytic activity.

© 2013 Elsevier Ltd. All rights reserved.

The *N*-aryl heterocycles are an important class of nitrogen containing compounds, that play a vital role in natural products, agrochemicals, material, and pharmaceutical products.¹ The *N*-aryl indoles are of much interest as antiallergic,² antipsychotic agents,³ analgesic,⁴ antiestrogen,⁵ cyclooxygenase (COX)-1 inhibitors,⁶ COX-2 inhibitors,⁷ melatonin receptor MT1 agonists,⁸ herbicidal agents,⁹ and anti HIV-1 agents.¹⁰ Furthermore, they are also used as important building blocks in the synthesis of various biologically active compounds.¹¹ Hence the development of simple and efficient methodology for the synthesis of *N*-aryl indole has attracted much attention in the past decades. Traditionally, *N*-aryl indoles have been synthesized by Ullmann-type cross coupling reactions.¹² However, this method requires high temperature and the use of stoichiometric amounts of copper catalyst thereby limiting its applications. The alternative procedure includes the palladium catalyzed coupling of heterocycles with aryl halides under mild reaction conditions.¹³ However, the use of more toxic and expensive palladium catalyst limits the attractiveness of these transformations.


Subsequently, much progress has been made for *N*-arylation of indoles in Ullmann-type coupling to soften reaction conditions by employing the correct combinations of copper catalysts, ligands, bases, and other additives.¹⁴ However, most of these protocols require ligands thereby increasing the cost. In addition toxic and non recyclable reaction media (DMF, DMSO) make protocols non

greener. Recently, some researchers reported *N*-arylation of heterocycles in an aqueous medium because of its low cost, safety, and environmentally benign nature.¹⁵ However, these protocols have limitations such as low solubility of substrates and ligands in aqueous medium, microwave heating and limited substrate scope. Hence, to develop an economical and sustainable protocol for this transformation that operates under environmentally favorable condition is of great interest. The progress toward more greener and recyclable reaction media such as glycerol, PEG, and ionic liquid had recently been made in cross coupling reactions.¹⁶ Glycerol is one of the most economical and environmentally benign solvents and has triggered research in the development of new strategies for coupling reactions because of its versatile physicochemical properties such as high polarity, low flammability, high boiling point, biodegradability, and ability to dissolve both organic as well as inorganic compounds. With the increase in biodiesel manufacture world-wide, the market saturation of glycerol, a by-product of biodiesel is inevitably increasing.¹⁷ Here, we report mild and highly efficient methodology for *N*-arylation of indoles with aryl halides using a copper catalyst and glycerol as a green and recyclable solvent (Scheme 1).

Initially, to optimize the reaction condition indole (**1a**) and iodobenzene (**2a**) were chosen as a model substrate for the C–N coupling reaction by using CuI catalyst in glycerol at 120 °C. Under this condition, the *N*-aryl indole (**3a**) was obtained in 47% yield (Table 1, entry1). Recent studies show that the DMSO acts as an oxidant for metal species in cross coupling reaction.¹⁸ In order to improve the yield of *N*-aryl indole (**3a**), we performed the above reaction using 1 equiv of DMSO as additive. Interestingly *N*-aryl

* Corresponding author. Tel.: +91 22 33612601; fax: +91 22 33611020.

E-mail addresses: bhanage@yahoo.com, bm.bhanage@gmail.com, bm.bhanage@ictmumbai.edu.in (B.M. Bhanage).



Scheme 1. Copper catalyzed N-arylation of indoles with aryl halides.

Table 1
Screening reaction conditions for N-arylation of indole with iodobenzene^a



Entry	Catalyst	Base	Time (h)	Temp (°C)	Yield ^b (%)
1	CuI	K ₂ CO ₃	24	120	91, 47 ^c
2	CuCl	K ₂ CO ₃	24	120	79
3	CuBr	K ₂ CO ₃	24	120	75
4	CuCl ₂	K ₂ CO ₃	24	120	50
5	Cu ₂ O	K ₂ CO ₃	24	120	70
6	Cu(OAc) ₂ ·5H ₂ O	K ₂ CO ₃	24	120	53
7	CuO	K ₂ CO ₃	24	120	43
8	CuSO ₄ ·5H ₂ O	K ₂ CO ₃	24	120	49
9	—	K ₂ CO ₃	24	120	—
10	CuI	—	24	120	Trace
11	CuI	KOH	24	120	74
12	CuI	CS ₂ CO ₃	24	120	64
13	CuI	KO ^t Bu	24	120	62
14	CuI	Na ₃ PO ₄	24	120	43
15	CuI	Et ₃ N	24	120	49
16 ^d	CuI	K ₂ CO ₃	24	120	63
17	CuI	K ₂ CO ₃	24	110	72
18	CuI	K ₂ CO ₃	20	120	80

^a Reaction conditions: iodobenzene (1 mmol), indole (1.5 mmol), Cu catalyst (0.1 mmol), base (2 mmol), DMSO (1 mmol), glycerol (2 ml), 120 °C, 24 h.

^b GC yield.

^c Reaction carried out without DMSO.

^d CuI (0.05 mmol).

indole (**3a**) was obtained in 91% yield (Table 1, entry 1). Encouraged by this result, we screened various copper catalysts and bases for the model reaction (Table 1). It was observed that among the various screened copper catalysts, CuI gave the best yield of the desired product and hence was used for further studies (Table 1, entries 1–8), whereas no product formation was observed in the absence of copper catalyst (Table 1, entry 9). Subsequently, we screened various bases for this transformation and it was observed that the base plays an important role; in the absence of base very less product was formed (Table 1, entry 10). Among screened bases, K₂CO₃ exhibited higher activity (Table 1, entries 1 and 11–15). We have also screened various solvents such as DMSO, DMF, toluene, and water for this transformation. It was found that DMSO and DMF provided trace amount of the desired product and solvents like toluene and water were ineffective for N-arylation of indole. However, decreasing the catalyst loading from 10 mol % to 5 mol %, decreases yield of **3a** (Table 1, entry 16). In addition, the effect of reaction temperature and time was also investigated. It was observed that the yield of **3a** decreases with a decrease in

temperature (Table 1, entry 17). While decreasing the time from 24 h to 20 h, low yield of **3a** was observed indicating that 24 h was an optimum time required for completion of the reaction (Table 1, entry 18).

With these optimized reaction conditions in hand, we have studied the scope and limitation of developed protocol for N-arylation of various heterocycles. As illustrated in Table 2, the reaction of *N*-heterocycles with various halides proceeds efficiently to furnish a wide range of *N*-aryl heterocycles in good to excellent yield.²⁰ Firstly, we have screened various indole derivatives and it was observed that 3-methyl-1*H*-indole gave excellent yield of 3-methyl-1-phenyl-1*H*-indole (Table 2, entry 2), whereas 5-methoxy-, 5-bromo-1*H*-indole on coupling with iodobenzene afforded good yield of corresponding products (Table 2, entries 3 and 4). The prominent feature of developed protocol is that the bromo substituent at C-5 position of indole remained intact.

The indole with electron withdrawing substituent such as 5-nitro-1*H*-indole also provided acceptable yield (65%) of 5-nitro-1-phenyl-1*H*-indole (Table 2, entry 5). Interestingly, less reactive aryl halide such as bromobenzene also couples with indole derivatives furnishing *N*-aryl indoles in good to appreciable yields (Table 2, entries 6 and 7). The reaction of indole with 1-bromo-2-iodobenzene afforded the selective mono coupling 1-(2-bromophenyl)-1*H*-indole in good yield under optimized reaction condition (Table 2, entry 8). Next, electronically different halo derivatives with indole were also studied. The *ortho*-substituted 1-iodo-2-methoxybenzene afforded the 1-(2-methoxyphenyl)-1*H*-indole in 73% yield indicating no steric effect in the coupling reaction (Table 2, entry 9). Furthermore, good to excellent yields were achieved regardless of the electronic nature of the substituents on the aryl iodide, and no considerable electronic effects were observed for both *meta*- and *para*-substituted aryl iodides (Table 2, entries 10–12). We extended the substrate scope of the present protocol by replacing the indoles with other heterocycles. The treatment of pyrrole with aryl iodide afforded the corresponding 1-phenyl-1*H*-pyrrole in 52% yield (Table 2, entry 13). Further, various imidazole and pyrazole derivatives such as 1*H*-imidazole, 2-phenyl-1*H*-imidazole, and 3-phenyl-1*H*-pyrazole were coupled with aryl iodide furnishing the corresponding **3l**, **3m**, and **3n** in 67%, 64%, and 62% yields respectively (Table 2, entries 14–16).

In consideration of economical view of the developed methodology, reusability of the catalytic system was examined for N-arylation of (**2a**) with (**1a**) as model reaction after the extraction of product with diethyl ether. 1 mmol of fresh DMSO should have been complemented to the reaction system before the next run, as very less product was observed for next run without fresh addition of DMSO. The recovered glycerol layer containing copper catalyst accomplishes the respective transformation up to four times with iodobenzene and indole as a substrate (Table 3). In this

Download English Version:

<https://daneshyari.com/en/article/5264978>

Download Persian Version:

<https://daneshyari.com/article/5264978>

[Daneshyari.com](https://daneshyari.com)