



Ruthenium-catalyzed oxidation of alkynes to 1,2-diketones under room temperature and one-pot synthesis of quinoxalines

Yuan Xu, Xiaobing Wan*

Key Laboratory of Organic Synthesis of Jiangsu Province, College of Chemistry, Chemical Engineering, and Materials Science, Soochow University, 199 RenAi Road, Suzhou, Jiangsu 215123, China

ARTICLE INFO

Article history:

Received 19 September 2012

Revised 22 November 2012

Accepted 30 November 2012

Available online 5 December 2012

Keywords:

Ruthenium

Alkynes

Oxidation

1,2-Diketones

Quinoxalines

ABSTRACT

A ruthenium-catalyzed alkyne oxidation to 1,2-diketones using Oxone under room temperature is reported. Both substrate scope and mechanism were discussed. Notably, combination of the alkyne oxidation and condensation cyclization in one pot offers a very efficient and convenient entry into quinoxaline derivatives.

© 2012 Elsevier Ltd. All rights reserved.

Introduction

1,2-Diketones appear frequently as structural sub-units in molecules of biological and medicinal interest¹ and are versatile building blocks capable of undergoing a variety of chemical transformations, especially for the synthesis of heterocyclic compounds.² The direct oxidation of properly substituted alkynes, which are prepared via Sonogashira coupling, might be the most straightforward method to synthesize the 1,2-diketone derivatives. The stoichiometric oxidation of alkynes has been widely studied, which suffers from generation of large amounts of wastes. Recently, the development of catalytic methods for the oxidation of alkynes to 1,2-diketones has been an attractive research area in organic chemistry.³ In addition, C–C bond cleavage of 1,3-diketones represents another promising method for producing 1,2-diketones.⁴ Despite these great achievements, from the standpoint of environmental benignity and overall cost, pursuing more efficient and practical catalytic systems with improved turnover is paramount. As part of our ongoing project to investigate new oxidation methods for 1,2-diketone,^{3j,l,n} we attempted to develop a mild process catalyzed by ruthenium with low catalyst loadings.

Results and discussions


Recently, Yang et al. reported a Ru-catalyzed oxidative cleavage of alkynes to carboxylic acids using Oxone.^{5a} Inspired by this

* Corresponding author. Tel./fax: +86 512 65880334.

E-mail address: wanxb@suda.edu.cn (X. Wan).

work, we investigated the Ru-catalyzed oxidation of 1,2-diph- enylethyne **1a** for benzil **2a** using Oxone as the oxidant under room temperature. Sodium bicarbonate was used as the buffer to maintain the neutral conditions.⁵ To our delight, a small amount of benzil **2a** was formed in the catalytic system (Table 1, entry 11). Interestingly, the use of TEMPO as a cocatalyst enhanced the conversion in a remarkable manner, resulting in the product **2a** in excellent yield. The catalyst is capable of up to 4600 turnovers with the substrate **1a** (Table 1, entry 1). In sharp contrast with Yang's work, no carboxylic acid was detected in this transformation. Negligible product **2a** was generated in the absence of Ru catalyst (Table 1, entry 10). When other Ru catalysts were used, product **2a** was achieved in moderate yield (Table 1, entries 13 and 14). In addition to Oxone, various oxidants were also employed in this study. The results are shown in Table 1, which indicated that Oxone is the best oxidant for the transformation (Table 1, entries 15–18). The effect of the solvent was also dramatic. Among the various solvents examined, MeNO₂/H₂O was the most suitable for the transformation under the catalytic system. Replacement of MeNO₂/H₂O with other solvents leads to a drop in yield (Table 1, entries 2–9). Finally, when the transformation was scaled up to 10 mmol, product **2a** was still achieved in excellent yield (Table 1, entry 19).

With the optimized oxidation conditions available, we embarked on the investigation of the alkynes' scope. A series of alkynes with different substituents on the aromatic ring were examined (Table 2). To our delight, aromatic alkynes bearing both electron-rich and electron-poor functional groups were oxidized to the corresponding 1,2-diketones in moderate to excellent yields.

Table 1
Optimization of reaction conditions^a


| Entry | Catalyst | Cocatalyst | Oxidant | Solvent | Yield ^b |
|-------|--|------------|-------------------------------|-------------------------------------|--------------------|
| 1 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | MeNO ₂ /H ₂ O | 92% |
| 2 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | DMF/H ₂ O | <5% |
| 3 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | DME/H ₂ O | <5% |
| 4 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | EA/H ₂ O | <5% |
| 5 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | ^t BuOH/H ₂ O | 35% |
| 6 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | MeCN/H ₂ O | <5% |
| 7 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | Dioxane/H ₂ O | <5% |
| 8 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | Toluene/H ₂ O | <5% |
| 9 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | DCE/H ₂ O | <5% |
| 10 | — | TEMPO | Oxone | MeNO ₂ /H ₂ O | <5% |
| 11 | [Ru(cymene)Cl ₂] ₂ | — | Oxone | MeNO ₂ /H ₂ O | 26% |
| 12 | — | — | Oxone | MeNO ₂ /H ₂ O | N.O. ^c |
| 13 | RuCl ₂ (PPh ₃) ₃ | TEMPO | Oxone | MeNO ₂ /H ₂ O | 41% |
| 14 | RuCl ₃ | TEMPO | Oxone | MeNO ₂ /H ₂ O | 67% |
| 15 | [Ru(cymene)Cl ₂] ₂ | TEMPO | O ₂ | MeNO ₂ /H ₂ O | N.O. |
| 16 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Benzoquinone | MeNO ₂ /H ₂ O | N.O. |
| 17 | [Ru(cymene)Cl ₂] ₂ | TEMPO | PhI(OAc) ₂ | MeNO ₂ /H ₂ O | 40% |
| 18 | [Ru(cymene)Cl ₂] ₂ | TEMPO | H ₂ O ₂ | MeNO ₂ /H ₂ O | N.O. |
| 19 | [Ru(cymene)Cl ₂] ₂ | TEMPO | Oxone | MeNO ₂ /H ₂ O | 94% ^d |

^a All reactions were carried out in the scale of 0.2 mmol in 4.0 mL of nitromethane and 0.5 mL of water in the presence of 0.5 mmol NaHCO₃, 10 mol % TEMPO, and 0.02 mol % Ru catalyst using Oxone (330 mg, 5% active oxygen) under room temperature for 12 h unless noted otherwise.

^b Isolated yield.

^c Not observed.

^d 10 mmol substrate was used.

Table 2
Ru-catalyzed alkyne oxidation for 1,2-diketones^a

^b 50 °C.

^a All reactions were carried out in the scale of 0.2 mmol in 4.0 mL of nitromethane and 0.5 mL of water in the presence of 0.5 mmol NaHCO₃, 10 mol % TEMPO, and 0.02 mol % [Ru(cymene)Cl₂]₂ using Oxone (330 mg, 5% active oxygen) under room temperature for 12 h unless noted otherwise.

Download English Version:

<https://daneshyari.com/en/article/5265058>

Download Persian Version:

<https://daneshyari.com/article/5265058>

[Daneshyari.com](https://daneshyari.com)