



Catalyst free, three-component approach for unsymmetrical triarylmethanes (TRAMs)



Bhavani Shankar Chinta, Beeraiah Baire*

Department of Chemistry, Indian Institute of Technology Madras, Chennai 600036, Tamilnadu, India

ARTICLE INFO

Article history:

Received 12 September 2016

Revised 18 October 2016

Accepted 22 October 2016

Available online 24 October 2016

Keywords:

Grignard reagents

Catalyst free

Triarylmethanes

Three-component

Cascade strategy

ABSTRACT

An efficient catalyst free, three-component, cascade approach has been designed and developed for the rapid assembly of unsymmetrical triarylmethane framework. The strategy employed aromatic aldehydes, Grignard reagents, and neutral aromatics as reaction partners, under simple thermal conditions. This process affords very broad scope for all three partners as it is performed in the absence of any external additives.

© 2016 Elsevier Ltd. All rights reserved.

The development of synthetic methodologies for quick assembly of complex molecules with biological and material importance is of great interest from synthetic, environment and economical point of view. Therefore, recently tremendous efforts have been directed towards the discovery and development of efficient atom- and step-economical synthetic methods [1]. Multicomponent, tandem strategies will provide an opportunity to rapidly generate structural complexity from simple subunits. Reagent free multi-component reactions are highly environment friendly and observed very rarely in organic synthesis.

Triarylmethanes (TRAMs) have attracted much attention because of their unique structural and physical properties [2]. The triarylmethane motif has become one of the basic core structures for fluorescent molecules [3]. This substructure is also finding increased application in medicinal chemistry, such as antitubercular and anticancer agents [4] as well as potassium ion (K⁺) channel blockers [5]. Recently, a variety of fluorescent TRAMs have been applied in live cell imaging [6], and several TRAMs bearing metal binding sites are employed as selective sensors for metal ions [7]. Thus, it is becoming increasingly important to have efficient synthesis routes to symmetrical as well as unsymmetrical triarylmethanes (Fig. 1).

The Friedel–Crafts arylation, which involves the reaction between diarylmethanol or related derivatives and arenes in the presence of an acid catalyst is one of the most employed methods

for the synthesis of triarylmethanes [8,9]. Recently, transition metal-promoted coupling strategies have emerged as an alternative methods to provide access to structurally divergent triarylmethane frameworks [10]. But, both these strategies have a common feature among them i.e., the catalyst promoted condensation between a mono-aryl unit and a di-arylmethane derivative (Scheme 1).

Several reports [11] appeared on the synthesis of triarylmethanes from three mono-aryl units. But all these methods employs either acid catalysts or transition metal catalysts. Here in, we report for the first time an acid and transition metal free, three-component domino strategy for the rapid generation of unsymmetrical triarylmethanes by employing simple thermal conditions. According to our designed strategy, an aromatic aldehyde **1** (Ar¹-CHO) in the absence of any external reagents, would undergo a tandem, sequential nucleophilic addition by an aromatic carbanion reagent (Ar²-M, M = MgX or Li) **2** and a neutral aromatic nucleophile (Ar³-H) **3**, under thermal heating to generate the triarylmethane derivative **4**.

To test our hypothesis, initially we have treated 4-(N,N-dimethyl) benzaldehyde **5**, with PhMgBr (1.3 equiv.) in presence of N-Me-indole (2 equiv.) in THF (Scheme 2). To our delight, the expected TRAM **6** was formed in 88% yield after heating the reaction mixture at 80 °C for 14 h. Formation of **6** can be explained as a two stage cascade process (Scheme 2). First the Grignard addition to the aldehyde **5** to give the magnesium diarylcarbinoloxide **7**, followed by nucleophilic trapping of the in situ generated *p*-azaquinone-methide intermediate **8** by the aromatic nucleophile and re-aromatization

* Corresponding author.

E-mail address: beeru@iitm.ac.in (B. Baire).






Fig. 1. Representative triarylmethane based drugs, dyes and natural products.




Scheme 1. A) Known strategies (bi- and three-component) for the synthesis of TRAMs; and B) our catalyst free, three-component, cascade approach.



Scheme 2. A) Preliminary investigation on the designed three-component approach for the synthesis of triarylmethanes; and B) Possible mechanism. a) When, we employed PhLi in place of PhMgBr, the reaction was very slow and even after 50 h at 80 °C only 23% of TRAM 6 was isolated along with 32% of the alcohol 6' (see Supporting information for details).

of iminium ion 9 affords TRAM 7. Encouraged by this result, we next performed this transformation at various temperatures 55 °C and 100 °C, but none of them afforded better yields.

Next we performed a substrate scope study for various ArMgX reagents as well as for neutral aromatic nucleophiles (Scheme 3) keeping the aldehyde 5 as the primary substrate. Initially, diversely functionalized and substituted ArMgX reagents, such as 2-thienyl, 1-naphthyl, *p*-tolyl, *m*- and *p*-anisyl, *m*-CF₃-phenyl-, *p*-F-phenyl etc., were employed in combination with various substituted indoles, to quickly generate structurally divergent and complex unsymmetrical TRAM derivatives 10–17. In continuation, *N*-substituted anilines were also employed as neutral aromatic nucleophiles along with variety of aromatic Grignard reagents. All these nucleophiles very smoothly underwent the three-component



Scheme 3. Scope study for Grignard reagents and aromatic nucleophiles with *p*-(dimethylamino)benzaldehyde. a) reagents and conditions: aldehydes = 50 mg, RMgX (1.3 equiv.), Ar-H (2 equiv.); 80 °C; b) ~1 equivalent of excess Ar-H was recovered in all cases; c) yields after chromatographic purification; d) When, we employed 2-thienylli in place of 2-thienylMgBr, the reaction was very slow and even after 50 h at 80 °C only 13% of TRAM 18 was isolated along with 52% of the alcohol 18' (see Supporting information for details).

Download English Version:

<https://daneshyari.com/en/article/5265880>

Download Persian Version:

<https://daneshyari.com/article/5265880>

[Daneshyari.com](https://daneshyari.com)