



Aniline–terephthalaldehyde resin *p*-toluenesulfonic acid (ATRT) salt as efficient mild polymeric solid acid catalyst



Kiyoshi Tanemura*, Tsuneo Suzuki

School of Life Dentistry at Niigata, Nippon Dental University, Hamaura-cho, Niigata 951-8580, Japan

ARTICLE INFO

Article history:

Received 24 July 2013

Revised 24 September 2013

Accepted 30 September 2013

Available online 5 October 2013

Keywords:

Solid acids

Tetrahydropyranylation

Aniline–aldehyde resin

Deprotection

Silyl ethers

ABSTRACT

Aniline–terephthalaldehyde resin *p*-toluenesulfonic acid (ATRT) salt was easily prepared by the reaction of aniline with 1.25 equiv of terephthalaldehyde in the presence of 1.0 equiv of *p*-toluenesulfonic acid at 75 °C for 24 h in EtOH. ATRT efficiently catalyzed the tetrahydropyranylation of alcohols and deprotection of tetrahydropyranyl (THP), triethylsilyl (TES), and *tert*-butyldimethylsilyl (TBDMS) ethers. Deprotection of dodecyl THP ether and dodecyl TBDMS ether catalyzed by ATRT proceeded faster than those by pyridinium *p*-toluenesulfonate (PPTS). ATRT was reused without significant loss of activities.

© 2013 Elsevier Ltd. All rights reserved.

Use of solid acid catalysts in organic syntheses and industrial purposes provides some merits such as simple work-up procedure, repeated use of the catalyst, and less corrosion of the reaction vessel. A number of strongly acidic resin catalysts such as Amberlyst 15,¹ Dowex 50 series,² Nafion NR50,³ and Nafion SAC-13⁴ have been developed.^{5,6} These solid acids have limitations when applied to compounds involving acid-sensitive groups. On the other hand, a survey of literature revealed that commercially available weakly acidic resin catalysts seem to be limited to a few examples such as poly(4-vinylpyridinium *p*-toluenesulfonate) (polyPPTS)⁷ and Poly(4-vinylpyridinium chloride) (polyPCL).⁸ In addition, these catalysts are expensive and their frequency of use has been limited.

Aniline–aldehyde resins are one of the important synthetic polymers.⁹ Formaldehyde (FA), furfural, and benzaldehyde have been employed as aldehyde components. For example, aniline–formaldehyde resins have been prepared by the treatment of aniline (AN) with formalin in the presence of hydrochloric acid followed by neutralization with NaOH.¹⁰ They are superior to elasticity, impact resistance, and electric insulation, and have been employed for electrical insulators.⁹

We found that aniline–terephthalaldehyde resin *p*-toluenesulfonic acid (ATRT) salt acted as a mild polymeric solid acid catalyst. The resin can be synthesized from the condensation of AN with terephthalaldehyde (TPA) in the presence of *p*-toluenesulfonic acid (TsOH) by *only one step*. In this Letter, we wish to report the results for the synthesis of ATRT. In addition, tetrahydropyranylation of

alcohols and deprotection of tetrahydropyranyl (THP) and silyl ethers catalyzed by ATRT are described.

First, we prepared aniline–formaldehyde resin *p*-toluenesulfonic acid (AFRT) salt. AN was treated with various amounts of formalin in the presence of 1.0 equiv of TsOH at room temperature in water. Since the produced solids were soluble to hot DMSO and DMF, the resins were post-cured at 160 °C for 6 h. The results are summarized in Table 1. Acid density of AFRT was determined by acid titration. When the ratio of AN:FA is 1.0:1.5, tetrahydropyranylation of 1-dodecanol at room temperature for 8 h in CH₃CN gave dodecyl THP ether (**1b**) in 91% yield. However, AFRT was partially soluble in hot DMSO and DMF (entry 1).¹¹ When the ratio of AN:FA is 1.0:3.0, the yield of THP ether **1b** decreased (10%) although it was insoluble in hot DMSO and DMF (entry 4).

We examined the formation of aniline–benzaldehyde resin *p*-toluenesulfonic acid (ABRT) salt. However, the reaction of AN with 1.5 equiv of benzaldehyde in the presence of 1.0 equiv of TsOH at 75 °C for 24 h in EtOH did not give the solid.

Next, we conducted the synthesis of ATRT. AN was treated with various amounts of TPA in the presence of 1.0 equiv of TsOH at 75 °C for 24 h in EtOH. The results are shown in Table 2. Soft material was obtained from the case of AN:TPA = 1.0:0.75. When the ratio of AN:TPA is 1.0:1.0, THP ether **1b** was obtained in 97% yield after 2.5 h, but the obtained resin was somewhat wet (entry 1). When the ratio of AN:TPA is 1.0:1.25, THP ether **1b** was obtained in 98% yield after 2.5 h in CH₃CN (entry 2).¹² We employed ATRT (AN:TPA = 1.0:1.25) for the reactions described below. ATRT was quite insoluble to hot water and hot organic solvents such as hexane, benzene, CH₂Cl₂, CHCl₃, THF, acetone, CH₃CN, MeOH, DMSO,

* Corresponding author. Fax: +81 25 267 1134.

E-mail address: tanemura@ngt.ndu.ac.jp (K. Tanemura).

Table 1
Solubility and activity of AFRT

Entry	AN:FA	Solubility ^a		Acid density (mmol g ⁻¹)	THP ether 1b ^b	
		DMSO	DMF		Time (h)	Yield ^c (%)
1	1.0:1.5	++	++	0.70	8	91
2	1.0:2.0	+	+	0.59	8	15
3	1.0:2.5	+	+	0.43	8	11
4	1.0:3.0	–	–	0.30	8	10

^a Solubility to hot solvents:–(insoluble) < + < ++ (partially soluble).^b Reagents and conditions: 1-dodecanol (1.0 mmol), DHP (1.5 mmol), AFRT (200 mg), CH₃CN (5 mL), room temp.^c Isolated yields.**Table 2**
Solubility and activity of ATRT

Entry	AN:TPA	Solubility ^a		Acid density (mmol g ⁻¹)	THP ether 1b ^b	
		DMSO	DMF		Time (h)	Yield ^c (%)
1	1.0:1.0	–	–	1.70	2.5	97
2	1.0:1.25	–	–	1.35	2.5	98
3	1.0:1.5	–	–	1.05	3	94
4	1.0:2.0	–	–	0.65	3	100
5	1.0:2.5	–	–	0.24	2.5	95






^a Solubility to hot solvents:–(insoluble) < + < ++ (partially soluble).^b Reagents and conditions: 1-dodecanol (1.0 mmol), DHP (1.5 mmol), ATRT (200 mg), CH₃CN (5 mL), room temp.^c Isolated yields.

and DMF. ATRT possessed the composition of C_{1.000}H_{0.875}N_{0.053}-O_{0.161}S_{0.017}, indicating that the amount of salt formation was 32% per aniline unit.

In order to increase the acid density, further treatment of ATRT with TsOH at 90 °C for 24 h in water was carried out. The acid density increased from 1.35 to 1.92 mmol g⁻¹, but the activity of ATRT for tetrahydropyranylation of 1-dodecanol decreased (6 h, 97%). The reason is not clear at the present moment.

The results of tetrahydropyranylation of alcohols and deprotection of THP ethers catalyzed by ATRT are shown in Table 3. Various alcohols involving acetals, allylic hydroxy, triple bonds, and conjugated ketones were treated with 3,4-dihydro-2H-pyran (DHP) in CH₃CN to give the corresponding THP ethers in good to excellent yields. Similarly, various THP ethers were transformed to their corresponding alcohols in MeOH in good to excellent yields.

Table 3
Tetrahydropyranylation of alcohols and deprotection of THP ethers catalyzed by ATRT

Entry	Alcohol	Protection ^a		Deprotection ^b	
		Time (h)	Yield (%)	Time (h)	Yield (%)
1	CH ₃ (CH ₂) ₁₁ OH 1a	2.5	98	3.5	100
2	 2a	7	91	8	– ^c
3	 3a	8	100	7	90
4	PhCH ₂ OH 4a	2	94	3	87
5	 5a	24	88	3	81
6	 6a	24	100 ^d	8	92
7	 7a	15	90 ^e	7	97 ^f

^a Reagents and conditions: alcohol (1.0 mmol), DHP (1.5 mmol), ATRT (200 mg), CH₃CN (5 mL), room temp.^b Reagents and conditions: THP ether (1.0 mmol), ATRT (200 mg), MeOH (5 mL), room temp.^c Glycerol was obtained in 91% yield.^d DHP (2.0 mmol), ATRT (400 mg), CH₃CN (10 mL) were used.^e CH₂Cl₂ (5 mL) was used.^f MeOH (5 mL)–CH₂Cl₂ (5 mL) was used.

Download English Version:

<https://daneshyari.com/en/article/5271565>

Download Persian Version:

<https://daneshyari.com/article/5271565>

[Daneshyari.com](https://daneshyari.com)