



Basic ionic liquid promoted heterocyclization to access fused imidazopyridines



I. R. Siddiqui*, Shireen, Shayna Shamim, Malik Abdul Waseem, Afaf A. H. Abumhdi, Arjita Srivastava, Anjali Srivastava

Laboratory of Green Synthesis, Department of Chemistry, University of Allahabad, Allahabad 211002, India

ARTICLE INFO

Article history:

Received 23 May 2013

Revised 4 July 2013

Accepted 7 July 2013

Available online 15 July 2013

Keywords:

Ionic liquid

Imidazopyridine

Heterocyclization

Room temperature

Mild reaction conditions

ABSTRACT

A single step access to fused imidazopyridine involving [bmlm]OH promoted click cyclocondensation of *N*-methylisatin with 2-aminopyridine has been achieved. The title heterocyclic scaffolds were obtained in excellent yield with high purity. [bmlm]OH was found to be more attractive as it plays the role of solvent, base, and catalyst.

© 2013 Elsevier Ltd. All rights reserved.

Nitrogen-containing heteroaromatics and their analogous are pharmaceutically attractive scaffolds and are broadly present in naturally occurring and synthetic biologically active molecules. Fused imidazo[1,2-*a*]heterocycles such as benzimidazo[2,1-*a*]isoquinolines (**I**)¹ and pyrido[1,2-*c*]purines (**II**) are anticancer agents², pyridino[1,2-*a*]imidazo[5,4-*b*]indol (**III**)(Fig. 1) and dimeric analogues of 2 aminodipyrido[1,2-*a*:3',2'-*d*]imidazole, are potent anti-hypertensive compounds.^{3,4} They possibly interact with DNA and display biological and pharmaceutical activity.⁵ Moreover, drugs like zolpidem which is clinically used for the treatment of insomnia, olprinone which is used for the treatment of acute heart failure, minodronic acid for the treatment of osteoporosis, and zolimidine as an anti-ulcer agent (Fig. 2) have imidazo[1,2-*a*]pyridine as the core structure.⁶

A number of synthetic routes have been reported for the synthesis of imidazo[1,2-*a*]pyridine.⁷ The synthesis of imidazopyridines containing pyrrole as an auxiliary group also suffered limitations like scope, generality, substrate availability, hazardous catalysts, carcinogenic VOLs (volatile organic liquids), and unavoidable need of bases. Furthermore these synthetic protocols involved time consuming sequential multistep processes with harsh reaction conditions. Because of strong requirement of fused imidazo[1,2-*a*]pyridine scaffolds in medicament, there is a demand for developing a straightforward, simple, high yielding, and conve-

nient facile methodology for the synthesis of imidazo[1,2-*a*]pyridine from simple and readily available precursors.

The pressing need is the use of alternative reaction medium that circumvents the problems associated with many traditional organic solvents. Operations resulting in the construction of highly functionalized heterocycles remain a great scientific challenge. This has prompted us to utilize [bmlm]OH in intermolecular cyclization to facilitate the synthesis of fused imidazopyridine. Buoyed from the green credentials of task specific basic ionic liquids (TSBIL)^{8–10} as well as part of our ongoing endeavors for the development of simple, efficient, and versatile synthetic methodologies for the synthesis of biodynamic heterocyclic scaffolds,^{11–16} we report herein a single step methodology using 2-aminopyridine and *N*-methylisatin as reactants catalyzed by [bmlm]OH (1-butyl-3-methylimidazolium hydroxide) to furnish fused imidazopyridine scaffolds (Scheme 1).¹⁷

We have successfully developed an efficient method to prepare a number of fused imidazopyridine derivatives via basic ionic liquid promoted cyclocondensation of *N*-methylisatin and 2-aminopyridine at room temperature. The selection of base, catalyst, and solvent system is a main issue as per importance of environmental purity. However, in the previously reported work, catalysts like Cu, Pd, Rh, solvents such as DMF, ethanol, toluene, and bases like NaOH have already been used in standard reaction conditions. We have replaced toxic catalysts, hazardous bases, and carcinogenic volatile organic solvents with recyclable and non hazardous [bmlm]OH and achieved better yield of the product (89%).

* Corresponding author. Tel.: +91 532 2461236; fax: +91 532 2461157.

E-mail address: dr.irsiddiqui@gmail.com (I.R. Siddiqui).






Figure 1.



Scheme 1. Basic ionic liquid catalyzed synthesis of fused imidazo-pyridine.

Table 1
Influence of various solvents on the synthesis of **4a** using NaOH as base and CuI as catalyst^a



| Entry | Solvent | Base (10%) | Catalyst (10%) | Time (h) | Yield ^b (%) |
|-------|------------------|------------|----------------|----------|------------------------|
| 1 | H ₂ O | NaOH | CuI | 7 | 21 |
| 2 | Ethanol | NaOH | CuI | 5 | 37 |
| 3 | DCM | NaOH | CuI | 4 | 53 |
| 4 | THF | NaOH | CuI | 5 | 43 |
| 5 | DMF | NaOH | CuI | 6 | 48 |

^a Reaction conditions: *N*-methylisatin (2.0 mmol), 2-aminopyridine (2.0 mmol).

^b Isolated yields.


In order to explore the generality and feasibility of the strategy we attempted to investigate the optimization of reaction conditions using 2-aminopyridine and *N*-methyl isatin as reaction reactants. Various solvents were used in the presence of NaOH as a base and CuI as a catalyst to afford desirable products and the results are summarized in Table 1.

It was inferred from Table 1 that the best solvent in terms of product yield and reaction time was DCM (Table 1, entry 3) while ethanol, THF, CH₂Cl₂, and H₂O proved less effective as lower yield of product was obtained and longer reaction time was needed (Table 1, entries 2, 4, and 5). To achieve high product conversion the reaction was carried out in DCM with different catalysts like CuI, CuBr, ZnCl₂, and InCl₃ and it was found that CuI turns out to be the best catalyst for the transformation of reactants into products (Table 2, entry 1).

Reports regarding heterocyclization reactions in ionic liquid encouraged us to perform the reaction in this versatile reaction medium which in most of the cases acted also as a promoter and additionally sometimes as base. So this reaction involving model reactants was carried using ionic liquids [bmIm]OH and [bmIm]Br without inorganic base, organic solvent, and metal halide as catalyst and the results are summarized in Table 3. From Table 3 [bmIm]OH was found to be the most efficient as there was no need to use additional catalyst and base. Further the suitable work-up process for the reaction is an additional advantage of the protocol.

The effect of mol % of [bmIm]OH on product yield and reaction time was also optimized for the synthesis of fused imidazopyridine by using different mol % of the ionic liquid (Table 4). From the results shown in Table 4, entry 1 it was inferred that 20 mol % of [bmIm]OH proved to be the optimum catalyst loading for intermo-

Table 2
Influence of catalyst on the synthesis of **4a**^a




| Entry | Catalyst (10%) | Solvent | Catalyst (10%) | Time (h) | Yield ^b (%) |
|-------|-------------------|---------|----------------|----------|------------------------|
| 1 | CuI | DCM | NaOH | 4 | 63 |
| 2 | CuBr | DCM | NaOH | 5 | 45 |
| 3 | ZnCl ₂ | DCM | NaOH | 6 | 38 |
| 4 | InCl ₃ | DCM | NaOH | 5 | 46 |

^a Reaction conditions: *N*-methylisatin (2.0 mmol), 2-aminopyridine (2.0 mmol).

^b Isolated yields.

Table 3
Influence of Ionic liquid on the synthesis of **4a**^a




| Entry | Ionic liquid | Time (h) | Yield ^b (%) |
|-------|--------------|----------|------------------------|
| 1 | [bmIm]OH | 3 | 86 |
| 2 | [bmIm]Br | 6 | 71 |

^a Reaction conditions: *N*-methylisatin (2.0 mmol), 2-aminopyridine (2.0 mmol).

^b Isolated yields.

Table 4
Influence of different mol (%) of [bmIm]OH on **4a**^a



| Entry | [bmIm]OH (mol %) | Time (h) | Temp (°C) | Yield ^b (%) |
|-------|------------------|----------|-----------|------------------------|
| 1 | 20 | 3 | rt | 86 |
| 2 | 15 | 3 | rt | 71 |
| 3 | 25 | 3 | rt | 86 |
| 4 | 20 | 4 | rt | 86 |
| 5 | 20 | 2 | rt | 78 |
| 6 | 20 | 3 | 60 | 86 |

^a Reaction conditions: *N*-methylisatin (2.0 mmol), 2-aminopyridine (2.0 mmol).

^b Isolated yields.

lecular heterocyclization with the product yield of 86%. By decreasing the amount of catalyst to 15 from 20 mol % relative to substrate, the yield of product **4a** decreased (Table 4, entry 2) but by increasing the amount of catalyst from 20 to 25 mol %, no appreciable change in the yield of product was observed (Table 4, entry 3). It was also observed that when [bmIm]OH was used as solvent and catalyst, at higher reaction temperature no appreciable effect on the yield was observed (Table 4, entry 6).

The task specific basic ionic liquid was synthesized according to a reported procedure.¹⁸ Ionic liquid [bmIm]OH was recovered completely and recycled twice for the synthesis with no appreciable decrease in the efficiency of the process (Fig. 3). The procedure for the synthesis of [bmIm]OH, and procedure for its recycling are given in the experimental section (Fig. 3).¹⁷

The formation of **4a** may be rationalized by the condensation of the carbonyl group of *N*-methyl isatin with 2-aminopyridine which on cyclization gives imidazopyridine intermediate **3**. This intermediate undergoes tautomerization and dehydration under the reac-

Download English Version:

<https://daneshyari.com/en/article/5272404>

Download Persian Version:

<https://daneshyari.com/article/5272404>

[Daneshyari.com](https://daneshyari.com)