



Diterpenoids from *Isodon pharicus*

Yong Zhao^{a,b}, Sheng-Xiong Huang^a, Wei-Lie Xiao^a, Li-Sheng Ding^c, Jian-Xin Pu^a, Xian Li^a,
Li-Bin Yang^a, Han-Dong Sun^{a,*}

^aState Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, PR China

^bCollege of Chemistry and Engineering, Yunnan Normal University, Kunming 650092, PR China

^cChengdu Institute of Biology, Chinese Academy of Sciences, Chengdu 610041, PR China

ARTICLE INFO

Article history:

Received 10 December 2008

Revised 10 February 2009

Accepted 13 February 2009

Available online 20 February 2009

Keywords:

Isodon pharicus

Labiatae

Bispseurata F

Pharicinins A–C

NMR data

Cytotoxicity

ABSTRACT

A phytochemical investigation of *Isodon pharicus* led to the isolation of a novel asymmetric *ent*-kauranoid dimer, bispseurata F (**1**), and three new diterpenoids, pharicinins A–C (**2–4**). Their structures were elucidated by extensive spectroscopic analysis. Compound **1** features a unique linkage pattern of C-17 with C-11' to connect the two monomers. A possible biogenetic pathway of **1** was also proposed. Compounds **3** and **4** exhibited moderate inhibitory activity against NB4 and SH-SY5Y cell lines.

© 2009 Elsevier Ltd. All rights reserved.


Isodon species are well known to contain structurally diverse and bioactive diterpenoids.¹ Over the past 30 years, more than 50 *Isodon* species have been phytochemically investigated, and a large number of new *ent*-kauranoids, including dimeric diterpenoids with six kinds of linkage pattern, have been found by our group.^{2–7} However, the dimeric diterpenoid, formed by the Michael addition reaction and connected by the C-17–C-11', has never been reported.

Isodon pharicus (Prain) Hara, mainly distributed in the northwest of Sichuan Province and the southern district of Tibet Autonomous region, People's Republic of China, has been used for killing roundworm and treatment of inflammation of the eyes.^{8a} Previous studies on this plant resulted in the isolation of six *ent*-kauranoids, including a dimeric diterpenoid.^{8b–d} In our search for biologically active secondary metabolites from *Isodon* genus, a novel dimeric *ent*-kauranoid, bispseurata F (**1**), and three new diterpenoids, pharicinins A–C (**2–4**), were isolated from the aerial parts of *I. pharicus*, collected in Lhasa Prefecture, Tibet Autonomous region. Compound **1** was characterized with a unique C-17–C-11' connection pattern to link the two monomers together, which was proposed to be formed by the Michael addition reaction. In this Letter, we present the isolation and structure elucidation of these new diterpenoids and their cytotoxicity evaluation, as well as the hypothetically biogenetic pathway of **1**.

The aerial parts of *I. pharicus* (7.0 kg) were extracted with 70% aqueous acetone (3 × 40 L) at room temperature overnight. The extract was partitioned between EtOAc and H₂O, and an EtOAc extract (273 g) was chromatographed over a silica gel, eluted in a step gradient manner with CHCl₃–CH₃COCH₃ (1:0 to 0:1) to afford fractions A–F. Fraction C (12 g) was submitted to chromatography over a silica gel (petroleum ether–acetone, from 9:1 to 1:1), to give fractions C1–C4. Fraction C2 (850 mg) was purified by repeated chromatography over a silica gel (petroleum ether–acetone, from 40:1 to 1:1) and RP-18 column (32% MeOH–H₂O), to yield compound **1** (10 mg). Fraction B (42 g) was submitted to chromatography over a silica gel (petroleum ether–acetone, from 99:1 to 1:1), to give fractions B1–B5. Compounds **2** (23 mg) and **4** (5 mg) were acquired from fraction B2 (1.2 g), and compound **3** (7 mg) was obtained from fraction B4 (3.8 g) by RP-18 (40% MeOH–H₂O).

Bispseurata F (**1**), isolated as a white, amorphous powder, exhibited the molecular formula C₄₄H₆₀O₁₄ on the basis of the positive HRESIMS (*m/z* 835.3900 [M+Na]⁺, calcd 835.3880), corresponding to 15 degrees of unsaturation.⁹ The IR absorptions at 3430, 1731, and 1643 cm^{−1} indicated the presence of hydroxyl, carbonyl, and α,β-unsaturated ketone groups, respectively. The ¹H, ¹³C, and DEPT NMR spectra (Table 1) displayed 44 carbon resonances due to eight methyls (of which two belonged to acetyls), seven methylenes (including one olefinic methylene), sixteen methines (of which eight were oxygenated), and thirteen quaternary carbons (including one olefinic carbon and six carbonyls). Above evidences showed that **1** could be a dimeric diterpenoid.

* Corresponding author. Tel.: +86 871 5223251; fax: +86 871 5216343.
E-mail address: hdsun@mail.kib.ac.cn (H.-D. Sun).



Comparison of the NMR data of **1** with those of pseurata F¹⁰, the monomer of **1**, also isolated from this plant, revealed that the two diterpenoid units of **1** were **1a** and **1b** (Fig. 1). The ¹H and ¹³C NMR spectra of both **1a** and **1b** resembled those of pseurata F except that the α,β-unsaturated ketone in pseurata F was replaced by an isolated ketone [δ_C 218.7 (s, C-15)], a methine [δ_H 3.20, m (H-16), δ_C 50.3 (d, C-16)], and a methylene [δ_H 1.12, m (H₂-17), δ_C 35.7 (t, C-17)] in **1a**, while a methine [δ_H 4.05, m (H-11'), δ_C 48.9 (d, C-11')] in **1b** substituted for a methylene [δ_C 38.7 (t, C-11)] in pseurata F. All these key changes of the characteristic signals were suggestive that the subunits **1a** and **1b** are connected by a single carbon-carbon bond (C17–C11'), which was proven by the ¹H–¹H COSY correlations of H-14/H-13/H-16/H₂-17/H-11'/H-9', as well as the observed HMBC correlations from H₂-17 to C-13 (δ_C 60.4), C-15 (δ_C 218.7), C-16 (δ_C 50.3), C-9' (δ_C 53.0), and C-12' (δ_C 209.0) (Fig. 1). In addition, since the correlations observed in the HMBC spectrum of subunits **1a** and **1b** were consistent with those of pseurata F, this confirmed the deduction above.

The relative configuration of compound **1** was established on the basis of the ROESY correlations of H-11' with H-13 α , H-16, and Me-20', of H-17 α with H-9' β , and of H-13 α with H-14 α as shown in the computer-generated 3D drawing (Fig. 1). In addition, the other correlations observed in the HMBC and ROESY spectra of **1** were in good agreement with those of pseurata F, this confirmed that the compound **1** was bispseurata F as shown.

Pharicin A (**2**) was obtained as a white powder, and its molecular formula C₂₂H₃₄O₃ was determined by the positive HRESIMS (m/z 369.2404 [M+Na]⁺, calcd 369.2405), with 6 degrees of unsaturation.¹¹ Apart from the resonances of an acetyl, twenty carbon signals corresponding to three tertiary methyls, eight methylenes (including one oxygenated and one sp² methylene), five methines (including one oxygenated and two sp² methines), and four quaternary carbons (including one sp² carbon) were observed in the NMR spectra, which were characteristic signals of an isopimarane diterpenoid. Considering from the biogenetic pathway and the fact that compound **2** was isolated from *Isodon* genus, compound **2** should be an *ent*-isopimarane diterpenoid.

The HMBC correlations from H-14 (δ_H 4.02) to C-7 (δ_C 119.0), C-12 (δ_C 35.5), C-15 (δ_C 149.6), and C-17 (δ_C 15.9) and from H₂-18 (δ_H 3.80 and 3.92) to C-3 (δ_C 36.5), C-5 (δ_C 44.7), C-19 (δ_C 18.2), and OAc (δ_C 170.9) suggested that a hydroxyl group and an acetoxy group were attached to C-14 and C-18, respectively. The HMBC correlations of H-15 (δ_H 6.30) with C-12 (δ_C 35.5), C-13 (δ_C 43.1), C-14 (δ_C 77.3), and C-17 (δ_C 15.9), of H₂-16 (5.10 and 5.19) with

C-13 (δ_C 43.1), C-14 (δ_C 77.3), and C-17 (δ_C 15.9), and of H-7 (δ_H 6.33) with C-6 (δ_C 23.6), C-9 (δ_C 51.9), and C-14 (δ_C 77.3) indicated that the two double bonds were located at C7–C8 and C15–C16, as shown in Figure 2. In the ROESY spectrum of **2**, H-14 correlated to H-9 β and H-11 β , and Me-17 correlated to H-11 α . These observations indicated that H-14 adopt a β -orientation, while Me-17 is α -oriented. Therefore, **2** was elucidated as 14 α -hydroxy-18-acetoxy-*ent*-isopimar-7,15-diene.

The molecular formula (C₂₂H₃₂O₆) of pharicin B (**3**)¹² was deduced from its HRESIMS (m/z 415.2089 [M+Na]⁺). The ¹H and ¹³C NMR data revealed that compound **3** resembled pseurata C¹³ with the presence of a hydroxyl at C-12 in **3** rather than a carbonyl at the same position in pseurata C, which was supported by the HMBC correlations from H-12 to C-9, C-13, C-14, and C-16. The HO-12 was assigned as the α -oriented on the basis of the ROESY correlations of H-12/ H-17 α (Fig. 3). Therefore, the structure of **3** was determined as 3 β -acetoxy-7 α ,12 α ,14 β -trihydroxy-*ent*-kaur-16-en-15-one.

The ¹H and ¹³C NMR data of pharicin C (**4**)¹⁴ were similar to those of leukamenin E¹⁵, and the only difference was that an acetoxy was located at C-19 in **4**, which was proven by the HMBC correlations from H₂-19 (δ_H 4.41 and 4.16) to OAc (δ_C 170.9). The relative configuration of **4** was identical to that of leukamenin E according to the ROESY correlations. Consequently, **4** was characterized as 3 β ,19-diacetoxy-7 α ,14 β -dihydroxy-*ent*-kaur-16-en-15-one.

Bispseurata F (**1**) was the first example of dimeric *ent*-kaurane diterpenoid connected by direct linkage of C-17 with C-11'. To clarify whether it is possible to produce **1** from pseurata F in the course of extraction or isolation by the Michael addition dimerization, we carried out some additional experiments, that is, mixed compound pseurata F with silica gel in different solutions (petroleum ether, CHCl₃, EtOAc, Me₂CO, and CH₃OH), kept the samples under air at room temperature for one month, and then monitored by the HPLC with **1** and pseurata F as controls. However, **1** was not detected in all mixtures. This showed that bispseurata F (**1**) is a naturally occurring product. A plausible biosynthetic pathway for **1** is proposed, as shown in Scheme 1. A Michael addition reaction is proposed to be the key step of this way. The dimerization of this type of diterpenoid dimer is worth further studies.

Compounds **1–4** were evaluated for their cytotoxicity against the NB4 (acute promyelocytic leukemia), A549 (lung cancer), PC-3 (prostate cancer), MCF-7 (breast cancer), and SH-SY5Y (neuroblastoma) human cell lines, using the sulforhodamine B (SRB)

Download English Version:

<https://daneshyari.com/en/article/5272464>

Download Persian Version:

<https://daneshyari.com/article/5272464>

[Daneshyari.com](https://daneshyari.com)