

Synthesis of a singly bridged resorcinarene-dimer by Mannich reaction with *N,N'*-diethylethylenediamine and formaldehyde: self-inclusion of bridging diazahexamethylene unit in methanol

Osamu Morikawa, Shingo Matsubara, Satoshi Fukuda, Kiyoshi Kawakami, Kazuhiro Kobayashi and Hisatoshi Konishi*

Department of Materials Science, Faculty of Engineering, Tottori University, 4-101 Koyama-minami, Tottori 680-8552, Japan

Received 27 July 2007; revised 11 August 2007; accepted 16 August 2007

Available online 21 August 2007

Abstract—Mannich reaction of resorcin[4]arene with *N,N'*-diethylethylenediamine and formaldehyde produced a singly bridged resorcinarene-dimer bearing sixteen hydroxyl groups at their peripheral positions, which exists in a closed capsular conformation including the diazahexamethylene bridge chain between the two resorcinarene units in CD₃OD.

© 2007 Elsevier Ltd. All rights reserved.

Calix[4]arenes and resorcin[4]arenes are [1₄]metacyclophane compounds, which can be readily prepared by a one-step procedure without using high dilution conditions.¹ These macrocycles have been used as building blocks for constructing molecular capsules.² A family of [1₄]metacyclophane-dimers carrying one intercavity linkage are attractive because the two bowl-shaped subunits arrange in an open-capsule or closed-capsule conformation as shown in Figure 1, and this conformational change may be achieved by varying the solvent polarity or encapsulating smaller molecules.

The synthesis of singly bridged dimers generally requires special precursors.³ They must be regioselectively func-





Figure 1. Conformational features in singly bridged resorcin[4]arene dimer.

Keywords: Self-inclusion; Capsule; Resorcin[4]arene dimer; Mannich reaction.

* Corresponding author. Tel.: +81 857 31 5262; fax: +81 857 31 5331; e-mail: konis@chem.tottori-u.ac.jp

tionalized to avoid bridging between two or more subunits. However, a few examples of the direct synthesis of dimers by the reaction of unmodified subunits with bifunctional reagents have been reported.⁴ The Mannich reaction of unmodified resorcinarene with ethylenediamine and an excess of formaldehyde yielded a carcer-and-type molecule in which two tetrabenzoxazine units are connected by four bridges.⁵ We are interested in extending this condensation to obtain a family of singly bridged resorcinarene dimers having sixteen hydroxyl groups.⁶ Such dimers might have interesting host properties, since strong electron-donating hydroxyl groups increase the electron density of the cavity. In these systems, cation- π and/or CH- π interactions between the electron-rich aromatic rings and guest molecules play an important role in stabilizing the inclusion complexes.⁷ We now report the facile synthesis of a singly bridged resorcin[4]arene dimer via the Mannich reaction of unprotected resorcinarene, *N,N'*-diethylethylenediamine and formaldehyde. Furthermore, we discuss its characteristic conformational properties in solution based on ¹H NMR spectroscopy.

The condensation of resorcin[4]arene **1**, *N,N'*-diethylethylenediamine **2** and formaldehyde in a 1:0.4:0.8 molar ratio was carried out in a 1:1 mixture of methanol and acetic acid at 50 °C for 5 h (Scheme 1). After removal of most of the solvent, the residual material



Scheme 1. Synthesis of a singly bridged resorcinarene dimer by the Mannich reaction.

was triturated with cold methanol to leave crude **3**, which was purified by crystallization from hot methanol to produce dimer **3** as an adduct with two molecules of acetic acid in 26% yield.[†] The Mannich reaction of resorcin[4]arenes can be carried out in ethanol–toluene.⁸ However, in this solvent system, the reaction with the diamine gave an intractable material. Furthermore, the reaction with piperazine produced a complex mixture from which we could not isolate any distinct product. It is presumed that the formation of the dimer significantly depends on the reaction conditions and the condensation agent.

The structure of dimer **3** was confirmed by elemental analysis, mass spectrometry and ¹H NMR spectroscopy. The ¹H NMR spectrum of dimer **3** in CD₃OD at 30 °C (Fig. 2A) shows five singlets at 6.12, 6.28, 7.24, 7.41 and 7.42 ppm in a 1:2:2:1:1 ratio for the aromatic protons, two sets of quartets at 4.49 and 4.50 ppm for the bridging methine protons and two sets of doublets at 1.70 and 1.71 ppm for the methyl protons. These data are in good agreement with C_s symmetrical structure of the mono-substituted resorcinarene **3**.

The chemical shifts of the bridging chain, H_a, H_b, H_c and H_d (see Fig. 2 for labels), are significantly affected by the solvent. In CD₃OD, dimer **3** displayed a triplet at –1.18 ppm and a quartet at 1.04 ppm for the *N*-ethyl groups (H_a and H_b). The central ethylene protons (H_c) and benzyl methylene protons (H_d) appear at 1.71 and 3.47 ppm, respectively. On the other hand, in DMSO-*d*₆, the ¹H NMR spectrum of **3** (Fig. 2B) shows H_a,




Figure 2. 400 MHz ¹H NMR spectra of **3**: measured in (A) CD₃OD; (B) DMSO-*d*₆. Asterisks (*) and crosses (x) indicate acetic acid and residual solvent protons, respectively.

H_b, H_c and H_d signals at 0.91, 2.40, 2.65 and 3.72 ppm, respectively. The chemical shift differences of those observed in CD₃OD and DMSO-*d*₆ for H_a, H_b, H_c and H_d are 2.09, 1.36, 0.94 and 0.25 ppm, respectively. These data suggest that, in methanol, the location of the protons of the bridging chain is influenced by the ring current shielding effects of the resorcinarene cavities.

The dimer is isolated as an adduct with two molecules of acetic acid, hence it is believed to exist as an ammonium salt. Therefore, the cation–π and/or CH–π interaction between the bridging chain and resorcinarene cavities is presumed to play a dominant role in the control of the conformation of the dimer since the resorcinarene anions strongly interact with ammonium ions. These interactions may occur in either an intra- or intermolecular fashion. To discriminate between these interactions, we examined the concentration dependence of the chemical shifts of **3** in CD₃OD. Figure 3 shows the high-field region of the ¹H NMR spectra of **3** as a function of the concentration. There are small, but clear changes in the chemical shift upon going from 0.27 to 13.5 mM. The upfield shifts of these protons with the decreasing concentration of **3** are not consistent with the intermolecular interaction, ruling out the possibility of the multi-component capsule formation. Thus, we speculate that **3** exists in a closed-capsule type conformation, in which the linkage moiety is included in the enclosed cavity, although the interaction in an intramolecular

[†] Preparation of dimer **3**: Resorcin[4]arene **1** (1.06 g, 2.0 mmol) was dissolved in hot methanol (20 ml). The solution was diluted with acetic acid (20 ml) and cooled to ambient temperature. To this were added *N,N'*-diethylethylenediamine (0.093 g, 0.8 mmol) and 36% aqueous formaldehyde (0.13 g, 1.6 mmol). The mixture was stirred for 2 h at ambient temperature and at 50 °C for 5 h. The solution was concentrated to one fifth under reduced pressure, and stored at 5 °C for 14 h to give solid material, which was washed with cold methanol to leave pure dimer **3** (380 mg) in 26% yield. mp 250 °C (dec). Anal. Calcd for C₇₂H₈₀O₁₆N₂(C₂H₄O₂)₂(H₂O)₆: C, 62.62; H, 6.92; N, 1.92. Found: C, 62.72; H, 6.55; N, 1.91. FAB-MS (MNBA) Calcd for C₇₂H₈₀O₁₆N₂: 1228.5. Found 1229.5 (M+1). δ_H (400 MHz; CD₃OD; Me₄Si; 30 °C) –1.18 (6H, br t, 6H, *N*-CH₂CH₃), 1.04 (4H, br q, *N*-CH₂CH₃), 1.70 (12H, d, bridge CH₃), 1.71 (12H, d, bridge CH₃), 1.71 (4H, s, CH₂CH₂), 1.96 (s, acetic acid), 3.47 (4H, s, Ar-CH₂-N), 4.49 (4H, q, bridge CH), 4.50 (4H, q, bridge CH), 6.12 (2H, s, ArH), 6.28 (4H, s, ArH), 7.24 (4H, s, ArH), 7.41 (2H, s, ArH), 7.42 (2H, s, ArH).

Download English Version:

<https://daneshyari.com/en/article/5275752>

Download Persian Version:

<https://daneshyari.com/article/5275752>

[Daneshyari.com](https://daneshyari.com)