

Diastereofacial selectivity in aldol-type condensation induced by optically pure α -sulfinyl acetate with α -substituted aldehydes

Claude Bauder

Institut de Chimie, UMR 7177 associé au CNRS—Université Louis Pasteur, 1 rue Blaise Pascal, 67000 Strasbourg, France

Received 4 January 2008; revised 6 February 2008; accepted 6 February 2008

Available online 10 February 2008

This Letter is dedicated with respect to the late Dr. Charles Mioskowski

Abstract

Highly diastereoselective aldol-type condensations induced by the chiral magnesium enolate of *tert*-butyl *p*-tolyl sulfinyl acetate were performed with optically pure alkyl or hydroxy α -substituted aldehydes. The addition of chiral α -sulfinyl acetate to various aldehydes, α -substituted or not, results mainly with a *syn* stereochemistry of the two newly created stereocenters. Applications were investigated for the synthesis of *syn* and *trans* 1,2-diol models and polysubstituted precursor for natural products.

© 2008 Elsevier Ltd. All rights reserved.

Keywords: Asymmetric aldolization; Chiral sulfinyl acetate; 1,2-Diol; β -Hydroxyester analogs

The control of the stereochemistry in the synthesis of complex natural products and their analogs requires various methodologies. In particular, much efforts have been devoted to the prevalent chiral enolate chemistry for the stereoselective construction of asymmetric centers in aldol condensation.¹ To this end, the use of chiral auxiliaries in asymmetric acetate aldol addition remains one of the most common and convenient strategy in organic synthesis. This procedure is widely developed in the literature to obtain with high diastereoselectivity, β -hydroxy acid derivatives which are chiral building blocks for the elaboration of bio-active compounds.²

For our part, we have been using for the last few years the chiral α -sulfinyl acetate in asymmetric aldol-type condensations³ as shown in Figure 1.

The well known optically active sulfoxide derivatives⁴ were first studied by Gilman, and then successively developed by Andersen and Solladié. They are prepared from diastereomerically pure (–)-(S) or (+)-(R) menthyl *p*-tolyl sulfinate **1** by treatment with a Grignard or other organo-





Fig. 1. General diastereoselective aldol-type reaction directed by optically pure α -sulfinyl ester.



Scheme 1. Synthesis of the enantiomers of *tert*-butyl *p*-tolyl sulfinyl acetate **2**.

metallic reagents.⁵ In this way, the two enantiomers of *t*-butyl *p*-tolyl sulfinyl acetate **2** (Scheme 1) were obtained

E-mail address: cbauder@chimie.u-strasbg.fr

in high yield from the appropriate menthyl sulfinate ester (–)-*S* **1** or (+)-*R* **1** and the lithium anion of *tert*-butyl acetate.

Subsequently, as shown in Figure 1, an aldehyde can be added now to the magnesium enolate of the α -sulfinyl acetate **2**, generating two new centers at C1 and C2 whose stereochemistry is governed by the chirality of the α -sulfinyl ester **2**. Such concept of asymmetric aldolization induced by optically pure sulfinyl ester was pioneered 30 years ago by Solladié and Mioskowski^{5f,6} and then has been used successfully in many applications in which an enantiomerically pure β -hydroxyester was needed.⁷ Since then, a number of examples have been reported involving the use of chiral α -sulfinyl acetamides,⁸ dihydrooxazoles,⁹ hydrazones¹⁰ or cyclic sulfinamides.¹¹ But in all these cases, the sulfoxide products were directly subjected to desulfurization resulting in β -hydroxycarbonyl derivatives (or β -aminoacid, when the sulfinyl ester **2** was added to an imine function¹²) with no consideration for the absolute configuration assignment at C1.

Recently, we have clearly assigned by X-ray analysis the absolute stereochemistry of the β -hydroxyester **3** (Fig. 2) which was obtained by direct aldol condensation from the magnesium enolate of the sulfinylester (–)-*S* **2** and achiral 3-benzyloxypropanal.³

Thus, these earlier results revealed a *syn* aldol relationship between the two newly created centers C1 and C2 as shown in molecule **3** (mp = 108 °C; $[\alpha]_D -199$ (*c* 0.85, CHCl₃)) with a coupling constant of 5.5 Hz between H-1 and H-2.¹³ We have also noticed an unexpected substituent effect when α,β -unsaturated aldehyde was used which necessarily involves a different transition state as previously postulated^{5e} (Fig. 3). On this basis we needed to explore the effect of α -substituted chiral aldehyde in the view to observe if in this case, the aldolization is matching either with achiral or α,β -unsaturated aldehyde situation.

In this Letter, we will examine a new series of condensations using various α -substituted (alkyl or hydroxyl) aldehydes. To our knowledge, no examples of α -substituted

aldehydes in aldol condensation with optically pure sulfinyl ester were described so far. We first achieved in good yield (65%) the condensation of (–)-*R* **2** and 3-benzyloxypropanal to isolate by simple crystallization product **4** (mp = 108 °C; $[\alpha]_D +195$ (*c* 1, CHCl₃)), which is the enantiomer of **3**. The structure as well as the *syn* stereochemistry of **4** were assigned by ¹H NMR analysis (*J* = 5.48 Hz between H-1 and H-2). That means a real transfer of chirality from the optically pure sulfinyl acetate to the prochiral α -unsubstituted aldehyde with high π -diastereofacial selectivity as shown in Figure 3.

Interested in the synthesis of natural products, we have selected compound **10** as a suitable fragment (Scheme 2). We envisaged our application in aldol condensation of the chiral sulfinylester (–)-*S* **2** with the readily available aldehyde **9** having an α -methyl group.

Aldehyde **9** was obtained by baker's yeast microbial reduction of ethyl acetylacetate to produce the (*S*)-ethyl β -hydroxy butanoate **5** in high enantiomeric purity.¹⁴ Then α -alkylation with LDA/MeI, according to the Frater–Seebach protocol¹⁵ led to the *anti* compound **6** in 63% yield. A sequential silylation–reduction of the ethyl (+)-(2*S*,3*S*) 3-hydroxy-2-methylbutyrate **6** followed by a Swern oxidation¹⁶ furnished the desired aldehyde **9** (69% yield for three steps). Subjection to an aldol condensation with the chiral sulfinylester (–)-*S* **2** provided the expected sulfoxide **10** as the sole isomer according to the ¹H NMR analysis of the crude material. Sulfoxide **10** was isolated by column chromatography from the unreacted sulfinyl ester, as an oily compound (68% yield, $[\alpha]_D -130$ (*c* 0.82, CHCl₃)). Its absolute stereochemistry was elucidated by NMR analysis. A coupling constant of 3 Hz is consistent with a *syn* stereochemistry between H-1 and H-2 in comparison with the value of 5.48 Hz for the known *syn* correlation in **3** and **4**. Moreover, an *anti* stereochemistry was clearly established (NOESY) between H-2 and H-3. Consequently, on the basis of ¹H NMR spectral data we assigned the configuration (*S*_S,1*R*,2*S*,3*R*,4*S*) to sulfoxide **10**.

Then, we extended this aldol-type condensation to α -hydroxy substituted aldehydes leading to the synthesis




Fig. 2. Configuration of **3** and **4** obtained by aldolization reaction.





Fig. 3. Illustration of a plausible chelated intermediate.



Scheme 2. Synthesis of sulfoxide **10**.

Download English Version:

<https://daneshyari.com/en/article/5278231>

Download Persian Version:

<https://daneshyari.com/article/5278231>

[Daneshyari.com](https://daneshyari.com)