

Highly efficient synthesis of α -amino amidines from ynamides by the Cu-catalyzed three-component coupling reactions

Ji Young Kim, Seok Hwan Kim, Sukbok Chang*

Center for Molecular Design and Synthesis (CMDs), Department of Chemistry and School of Molecular Science (BK21), Korea Advanced Institute of Science and Technology (KAIST), Daejeon 305-701, Republic of Korea

Received 16 December 2007; revised 13 January 2008; accepted 17 January 2008

Available online 20 January 2008

Abstract

Using ynamides as one of the reacting components, the Cu-catalyzed three-component reaction of sulfonyl or phosphoryl azides and amines affords α -amino amidines in high yields under mild conditions. Synthetic utility of the produced compounds was demonstrated in the diastereoselective alkylation of α -amino amidines bearing a chiral oxazolidinone moiety. It was also shown that α -amino imidates could be readily prepared by employing alcohols instead of amines.

© 2008 Elsevier Ltd. All rights reserved.


Keywords: α -Amino amidines; Ynamides; Three-component reactions; Copper catalysis

α -Amino amidines are widely found in natural products,¹ and used as important synthetic building units in the synthesis of iminopeptides² and antitumor or antibiotic compounds.³ However, only a few synthetic methods for the α -amino amidine compounds have been developed.⁴ For instance, Keung et al. presented a novel procedure for the synthesis of α -amino amidine via Sc-mediated Ugi condensation reactions.^{4c} We recently developed the three-component synthesis of *N*-sulfonylamidines by the copper-catalyzed coupling reaction of alkynes, sulfonyl azides, and amines.⁵ Imidates and amides were also prepared under the similar conditions with the use of alcohols⁶ or water⁷ instead of amines. More recently, phosphoryl azides were also found to be a facile reacting partner in the catalytic three-component reactions.⁸

Along with this line, we envisioned that α -amino amidines could be readily prepared by the Cu-catalyzed three-component coupling reactions when ynamides are employed as a source of terminal triple bonds. Hsung and Cintrat independently reported the highly efficient Cu-cata-

lyzed [3+2] cycloaddition reactions between ynamides and alkyl- or aryl azides producing 4-amino-1,2,3-triazoles.⁹ However, no example of utilizing ynamides in the catalytic three-component coupling reactions has been shown to afford α -amino amidines, which will be described herein (Scheme 1).

Since ynamides or ynamines bearing electron-deficient substituents have been shown to be highly useful synthetic precursors,^{10–12} several preparative routes are reported. For instance, Witulski paved the way for the synthesis of electronically tunable 1-alkynylamides using iodonium triflate salts in reaction with electron-deficient amines.¹³ Hsung demonstrated that *N*-alkynylation of oxazolidinones and lactams with alkynyl halides can be catalyzed by CuCN.¹⁴ Danheiser developed a modified procedure of *N*-alkynylation of carbamates, ureas, and sulfonamides



Scheme 1.


* Corresponding author. Tel.: +82 42 869 2841; fax: +82 42 869 2810.
E-mail address: sbchang@kaist.ac.kr (S. Chang).

using stoichiometric amounts of KHMDS and CuI.¹⁵ Urabe and Sato reported a route to *N*-(1-alkynyl)sulfonamides based on the Cu catalysis.¹⁶ Hsung's approach for the cross-coupling of amides with alkynyl bromides using catalytic amounts of CuSO₄ and 1,10-phenanthroline significantly broadened the availability of ynamides.¹⁷ Tam subsequently developed a method for the facile synthesis of acyclic ynamides by the slow addition of KHMDS to CuI catalytic systems.¹⁸

Following Tam's procedure with a slight modification, we were able to prepare a variety of ynamides in high yields substituted with several functional groups such as oxazolidinones, acyclic carbamates, and sulfonamides. A representative example is shown in Scheme 2. Cross-coupling of *N*-Boc-aniline and bromo(triisopropylsilyl)acetylene, which can be obtained by bromination of TIPS-acetylene with NBS,¹⁹ was successfully achieved by the action of CuI and 1,10-phenanthroline catalysts under the slow addition conditions.²⁰ Subsequent desilylation was readily carried out with tributylammonium fluoride, thus leading to the desired *N*-ethynyl-*N*-phenyl-(*tert*-butyl) carbamate (**1**) in 80% overall yield. Other substrates of ynamide derivatives could be readily prepared in similar yields using the same procedure.

With ynamides in our hands, we tried to optimize the reaction conditions for the synthesis of α -amino amidines using the Cu-catalyzed three-component reaction (Table 1). Initially, catalytic activity of various Cu species was examined in THF to reveal that CuI and CuCN were especially effective for the three-component reaction, leading to the desired product with high yields (entries 3 and 4, respectively). However, copper(II) complexes displayed reduced catalytic activities (entries 5–7). Among several solvents screened, chloroform turned out to be the best one with CuI catalyst (entry 9) while polar media were less effective (entries 11–13). Interestingly, no significant effects of external ligands were observed (entry 14).^{21,22}

Under the optimized conditions, a wide range of ynamides, azides, and amines were smoothly reacted to afford the corresponding α -amino amidine products in good yields (Table 2).²³ In addition to acyclic *N*-ethynyl carbamate (entry 1), cyclic analogs such as ynamides of oxazolidinone could also be readily employed as a viable type of substrates (entries 2–5). Interestingly, steric variation on the cyclic skeleton in these substrates does not alter efficiency of the coupling reactions. For instance, reactions with substrates bearing various substituents such as benzyl,



Scheme 2. Reagents and conditions: (a) *N*-Bromosuccinimide (1.2 equiv), AgNO₃ (0.1 equiv), acetone, 25 °C, 2 h; (b) *N*-Boc-aniline (0.8 equiv), CuI (0.25 equiv), 1,10-phenanthroline (0.3 equiv), KHMDS (1.1 equiv), toluene, 90 °C, 12 h; (c) TBAF (2.0 equiv), THF, 0 °C, 1 h.

Table 1

Optimization of the α -amino amidine synthesis^a

Entry	Catalyst	Solvent	Yield ^b (%)
1	CuCl	THF	35
2	CuBr·SMe ₂	THF	<1
3	CuI	THF	78
4	CuCN	THF	80
5	Cu(acac) ₂	THF	53
6	Cu(OAc) ₂	THF	59
7	Cu(OTf) ₂	THF	<1
8	CuI	CH ₂ Cl ₂	66
9	CuI	CHCl ₃	82
10	CuCN	CHCl ₃	66
11	CuI	DMF	43
12	CuI	CH ₃ CN	71
13	CuI	1,4-Dioxane	62
14 ^c	CuI	CHCl ₃	74

^a A mixture of *p*-toluenesulfonyl azide (0.6 mmol), **1** (0.5 mmol), diisopropylamine (0.6 mmol), and Cu catalyst (10 mol %) in solvent (1.0 mL) was stirred at 25 °C for 6 h.

^b H NMR yield (internal standard: 1,3-benzodioxole).

^c 20 Mol % of tris(benzyltriazolylmethyl)amine was added.

isopropyl, or *tert*-butyl group on the 4-position of oxazolidinone provided the corresponding products with high yields in all cases (entries 3–5, respectively). *N*-Sulfonamido acetylene was also employed as a substrate to afford α -sulfonamido amidine in a moderate yield (entry 6).

It was revealed that, like arenesulfonyl azides, alkylsulfonyl azides were also smoothly reacted leading to *N*-alkylsulfonyl amidines in high yields as demonstrated in entry 7. We were pleased to observe that phosphoryl azides displayed similar reactivity with ynamides to afford the corresponding α -amino-*N*-phosphoryl amidines in excellent yields (entries 8 and 9). It should be mentioned that the scope of amine species was very broad. Aniline derivatives as well as aliphatic amines were shown to be efficiently coupled in the presence of triethylamine (entry 10). Primary amines and amino esters were also readily employed in this coupling (entries 11 and 12, respectively).

The produced α -amino amidine compounds can be regarded as α -amino acid isosteres. For further synthetic applications, asymmetric alkylation of optically active α -amino amidines was next investigated (Table 3). It was especially interesting to see any steric influence of 4-substituents of the oxazolidinone skeleton on the diastereoselectivity during the alkylation process.²⁴ After examining various reaction conditions, it was found that deprotonation of the chiral α -amino amidines with LHMDS (in THF) followed by alkylation with electrophiles provided monoalkylated products in moderate to good chemical yields.

Not surprisingly, the observed diastereoselectivity turned out to be dependent on the steric environment on the oxazolidinone moiety despite the fact that the extent

Download English Version:

<https://daneshyari.com/en/article/5278449>

Download Persian Version:

<https://daneshyari.com/article/5278449>

[Daneshyari.com](https://daneshyari.com)