

Suzuki reaction of cyclopenta[*d*][1,2]oxazine in aqueous solvent with water-soluble phosphine ligand

Sung Yun Cho,* Seung Kyu Kang, Jin Hee Ahn, Jae Du Ha and Joong-Kwon Choi

Bio-Organic Science Division, Korea Research Institute of Chemical Technology, 100 Jang-dong, Yuseong-gu, Daejeon 305-600, Republic of Korea

Received 17 May 2006; accepted 24 May 2006

Available online 12 June 2006


Abstract—A convenient and facile modification of cyclopenta[*d*][1,2]oxazines through Suzuki cross-coupling reaction by use of water-soluble phosphine ligand was accomplished. In aqueous solvent, 7-iodocyclopenta[*d*][1,2]oxazines readily afford the corresponding 7-arylcyclopenta[*d*][1,2]oxazines in moderate to excellent yields.

© 2006 Elsevier Ltd. All rights reserved.

Cyclopenta[*d*][1,2]oxazine framework is seldom found in natural products or pharmaceuticals. There have been only a handful of reports on the synthesis of cyclopenta[*d*][1,2]oxazine and its derivatives, presumably due to limited stability of cyclopenta[*d*][1,2]oxazine skeletons.

Lloyd and Preston¹ first reported the synthesis of cyclopenta[*d*][1,2]oxazine by the reaction of diacylcyclopentadiene with hydroxylamine. They also reported that treatment of cyclopenta[*d*][1,2]oxazine with *N*-bromosuccinimide afforded 7-bromocyclopenta[*d*][1,2]oxazine. In an effort to pursue inhibitors of the protein tyrosine phosphatase 1B (PTP1B),² it was required to synthesize the targeting compounds via Suzuki cross-coupling of 7-iodocyclopenta[*d*][1,2]oxazines. Palladium catalyzed cross-coupling reaction of aryl halides with boronic acid is a powerful synthetic method for arylation in carbon–carbon bond formation.³ Aqueous-phase palladium catalyzed cross-coupling reactions are of interest with respect to environmental friendly synthetic process, the simplification of product separation and the economy of using water as a solvent in large-scale industrial syntheses.⁴ In particular, aqueous palladium catalyzed cross-coupling reactions were considered to be highly versatile and useful method for heat or base labile compounds for decomposition, such as cyclopenta[*d*][1,2]oxazines. Herein, we report a synthetic method for halogenated cyclo-

penta[*d*][1,2]oxazines and palladium-catalyzed cross-coupling reaction of 7-iodocyclopenta[*d*][1,2]oxazines by use of water soluble phosphine.




For initial studies, we examined the coupling of **1** and **2** in the presence of palladium acetate. All experiments were run at 50 °C and at room temperature with **1** and 2.0 equiv of lithium acetate, Cs₂CO₃, K₂CO₃ in CH₃CN–H₂O (1:1) or 1,4-dioxane. We failed to produce any biaryl adduct of cyclopenta[*d*][1,2]oxazine **3** without remaining substrate **1**, even under a condition where triethylamine acted as a base in DMF at room temperature. From this result, we found that cyclopenta[*d*][1,2]oxazines were not tolerable under heated condition. Indeed, we turned our attention to aqueous Suzuki aryl coupling of cyclopenta[*d*][1,2]oxazines with water soluble phosphine ligand mediated by palladium acetate. Initially, all of the 7-iodo and 7-bromocyclopenta[*d*][1,2]oxazines were synthesized through several steps (Table 1).

For iodination reaction, treatment of cyclopenta[*d*][1,2]oxazine by *N*-iodosuccinimide in MeCN in the presence of trifluoroacetic acid gave **1b** in excellent yield. Water soluble phosphine ligand was prepared by a known procedure as cited in the literature.⁵

Keywords: Suzuki reaction; Cyclopenta[*d*][1,2]oxazine; Water-soluble phosphine ligand.


* Corresponding author. Tel.: +82 42 860 7077; fax: +82 42 860 7160; e-mail: sycho@kriict.re.kr

Table 1.



Iodination reagent	Solvent	Yield (%)
ICI	AcOH	11
ICI	TFA/CH ₂ Cl ₂	80
NIS	TFA/MeCN	91

Table 2.



Entry	R'	Condition	Yield (%)
1	Br	CS ₂ CO ₃ , P(<i>o</i> -tolyl) ₃ , 1,4-dioxane, 50 °C	0 ^a
2	I	CS ₂ CO ₃ , P(<i>o</i> -tolyl) ₃ , 1,4-dioxane, 50 °C	0 ^a
3	Br	NaOAc, I , MeCN–H ₂ O (1:1)	0
4	I	NaOAc, I , MeCN–H ₂ O (1:1)	8
5	I	NaOAc, I , H ₂ O	7
6	I	CS ₂ CO ₃ , I , MeCN–H ₂ O (1:1)	8
7	I	NEt ₃ , I , MeCN–H ₂ O (1:1)	6
8	I	Toluene–MeOH–H ₂ O (1:1:1), I , CS ₂ CO ₃	66


^a No substrate was recovered.

For initial assessment of coupling reaction, Suzuki cross-coupling reactions were performed with a water

soluble phosphine ligand, [2-(di-*tert*-butyl phosphanyl) ethyl] trimethyl ammonium iodide **I**, as cited in the literature.⁶ The result given in Table 2 reveals that toluene was the solvent of choice to afford the desired 7-aryl-cyclopenta[*d*][1,2]oxazine along with CS₂CO₃ as a base without decomposition of the substrate at room temperature in moderate yield.

Suzuki reactions were not preceded under typical coupling condition with Pd(OAc)₂, tri(*o*-tolyl) phosphine in dioxane at 50 °C, and no substrate was recovered. (Table 2 entry 1 and 2) The reactions that were executed in water alone occurred more slowly than in water-miscible co-solvents (entry 5 in Table 2). With solvents, MeCN and water, the Suzuki reactions did not give rise to the desired product with satisfactory yields (entries 4–7). It was observed that the substrate was decomposed in either toluene or MeOH as a sole solvent without the addition of water. It was improved in yield with the addition of toluene in the reaction mixture, presumably because of better forming homogeneous reaction mixture (entry 8).

To evaluate the scope and limitation for this procedure to cyclopenta[*d*][1,2]oxazine acting as a substrate, we performed a number of coupling reaction with several aryl boronic acids and 7-iodocyclopenta[*d*][1,2]oxazines. All reactions were run at room temperature and cesium carbonate was used as a base of choice in toluene–MeOH–water (1:1:1) which was used as a solvent as given in Table 3. [2-(di-*tert*-butyl phosphanyl) ethyl] Trimethyl ammonium iodide **I** has allowed the first general coupling of aryl iodide to cyclopenta[*d*][1,2]oxazine at room temperature under nitrogen atmosphere. To

Table 3. Suzuki-coupling of 6-iodocyclopenta[*d*][1,2]oxazines

Compound no.	R ¹	R ²	Ar	Yield (%)
3a	Me	–CO ₂ Me (a)		65
3b	Me	a		75
3c	Me	a		65
3d	Me	a		67
3e	Me	a		75
3f	Me	a		78

Download English Version:

<https://daneshyari.com/en/article/5278798>

Download Persian Version:

<https://daneshyari.com/article/5278798>

[Daneshyari.com](https://daneshyari.com)