

# Cupric chloride promoted regioselective C-allylation of enaminones

Sarangthem Joychandra Singh, Okram Mukherjee Singh\*

Department of Chemistry, Manipur University, Canchipur, 795 003 Manipur, India

Received 8 October 2007; revised 8 April 2008; accepted 16 April 2008

Available online 20 April 2008

## Abstract

Regioselective allylation of enaminones using  $\text{CuCl}_2$  as the catalyst to give C-allylated products is reported for the first time. The C-allylated products undergo hydrolysis followed by a rearrangement yielding  $\beta$ -keto allyl enamides in excellent yields.  
© 2008 Elsevier Ltd. All rights reserved.


**Keywords:** Allylation; Enaminone;  $\beta$ -Keto allyl enamides; Ketene *N,S*-acetal

Cuprous and cupric salts are used as effective catalysts in several oxidative ring cyclization reactions,<sup>1</sup> and in the presence of oxygen and pyridine or amino ligands, they are considered as useful oxidizing systems for cleavage of hydrazides.<sup>2</sup> Another interesting application of copper salts is in the addition reaction of allylmetals to  $\text{C}=\text{Z}$  bonds ( $\text{Z}=\text{O}$  or  $\text{NR}$ ), which has emerged as a very useful carbon–carbon bond-forming reaction.<sup>3</sup> Metal-catalyzed allylic substitution is a useful process in organic synthesis for C–C and C–heteroatom bond forming reactions.<sup>4</sup> However, these approaches are still extremely limited in scope and functional group compatibility. Moreover, despite the potential utility of allylated  $\beta$ -keto amides as synthetic intermediates, their synthesis using copper salts as catalyst has been rarely studied.<sup>5</sup>

We reported some years ago the synthesis of biologically important 2,3-functionalized imidazo[1,2-*a*]pyridines via an unprecedented  $\text{CuCl}_2$ -induced oxidative ring closure of certain  $\alpha$ -oxoketene *N,S*-, *N,O*- and *N,N*-acetal intermediates.<sup>6</sup> Ketene *N,S*-acetals are highly versatile enamines widely used in the synthesis of heterocycles.<sup>7</sup> Junjappa and co-workers have recently reported the synthesis of substituted quinolines and quinoxalines via cyclization of ketene *N,S*-acetals under Vilsmeier–Haack conditions.<sup>8</sup>

We now report a facile synthesis of  $\beta$ -keto allyl enamides via regioselective allylation of *N,S*-acetals using cupric chloride as the catalyst.

Ketene *N,S*-acetals were prepared starting from ketene dithioacetals by displacement of the thiomethyl groups by the respective amines either in refluxing ethanol or in tetrahydrofuran.<sup>9</sup> Our various trial experiments to prepare either the C- or the N-allylated products of the ketene *N,S*-acetals, by treating **1a–l** with allyl bromide without using any catalyst in various solvents, resulted in failure. Thus refluxing *N,S*-acetals **1a–l** and allyl bromide in solvents such as ethanol, acetonitrile and tetrahydrofuran gave no significant results. Surprisingly, we observed the formation of  $\beta$ -keto enamides **2a–l** in excellent yields by treating the *N,S*-acetals and allyl bromide directly in the presence of cupric chloride and refluxing in THF for 2–3 h (Scheme 1).<sup>10</sup>



Scheme 1.

\* Corresponding author.


E-mail address: ok\_mukherjee@yahoo.co.in (O. M. Singh).

In order to evaluate the scope of this catalytic system, the range of metal salts was extended to various metal halides and acetates as shown in Table 1, guided by the template reaction between ketene *N,S*-acetal **1a** and allyl bromide. The CuCl<sub>2</sub>/THF combination was found to be the best affording the highest yield of 90% after refluxing for 3 h. Screening of different solvents revealed that THF was the most suitable. It was also observed that ZnCl<sub>2</sub> and SnCl<sub>2</sub> gave good yields of the product, while MgCl<sub>2</sub>, AlCl<sub>3</sub>, FeCl<sub>3</sub>, BiCl<sub>3</sub> and LaCl<sub>3</sub> gave poor yields of the desired products (Table 1). Moreover, it is noteworthy to mention that no C-allylation occurred in the absence of catalyst (Table 1, entry 15).

After optimizing the reaction conditions, we investigated the generality of this process. As can be seen from Table 2, various ketene *N,S*-acetals **1a–l** were C-allylated with allyl bromide to give the corresponding products in good to excellent yields.

We wanted to confirm that the reaction pathway occurred via C-allylation rather than N-allylation. For this study, we required N-allylated ketene *N,S*-acetals **3a–l** (Scheme 2) and it was anticipated that N-allylated adducts **3a–l** might undergo a 3-aza-Claisen rearrangement to give C-allylated products.<sup>11</sup> Recently, Oshima et al. reported<sup>12</sup> a facile method of synthesizing 2,2-disubstituted-4-pentenitriles via aza-Claisen rearrangement of *N*-allyl-*N*-(phenylethynyl)arenesulfonamides. Thus, as shown in Scheme 2, we treated *N,S*-acetals **1a–l** with allyl bromide at 0 °C to 25 °C in the presence of 2 equiv of sodium hydride in


Table 1  
Optimization of the reaction conditions for the C-allylation of **1a** with allyl bromide<sup>a</sup>



Entry	Catalyst	Solvent	Time (h)	Yield (%)
1	CuCl <sub>2</sub>	DMF	6	65
2	CuCl <sub>2</sub>	Benzene	10	60
3	CuCl <sub>2</sub>	Toluene	5	60
4	CuCl <sub>2</sub>	THF	3	90
5	CuBr <sub>2</sub>	THF	6	60
6	CuI <sub>2</sub>	THF	6	45
7	Cu(OAc) <sub>2</sub>	THF	5	45
8	MgCl <sub>2</sub>	Toluene	12	10
9	AlCl <sub>3</sub>	Toluene	12	20
10	FeCl <sub>3</sub>	Toluene	10	35
11	BiCl <sub>3</sub>	Toluene	10	15
12	LaCl <sub>3</sub>	Toluene	8	10
13	ZnCl <sub>2</sub>	THF	6	50
14	SnCl <sub>2</sub>	THF	6	50
15	None	THF	24	0


<sup>a</sup> *N,S*-Acetal (5.0 mmol), allyl bromide (5.1 mmol), catalyst (5.0 mmol), solvent (20–30 mL).

Table 2  
β-Keto allyl enamides **2a–l** produced via C-allylation of ketene *N,S*-acetals **1a–l**<sup>a</sup>



Entry	R <sup>1</sup>	R <sup>2</sup>	Product	Yield (%)	Mp (°C)
1	C <sub>6</sub> H <sub>5</sub>	CH <sub>3</sub>	<b>2a</b>	90	103
2	C <sub>6</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	<b>2b</b>	92	120
3	C <sub>6</sub> H <sub>5</sub>	C <sub>3</sub> H <sub>7</sub>	<b>2c</b>	86	105
4	C <sub>6</sub> H <sub>5</sub>	CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>	<b>2d</b>	85	120
5	4-MeC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	<b>2e</b>	90	122
6	4-MeC <sub>6</sub> H <sub>4</sub>	C <sub>2</sub> H <sub>5</sub>	<b>2f</b>	92	113
7	4-MeC <sub>6</sub> H <sub>4</sub>	C <sub>3</sub> H <sub>7</sub>	<b>2g</b>	90	107
8	4-MeC <sub>6</sub> H <sub>4</sub>	CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>	<b>2h</b>	80	125
9	4-MeOC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	<b>2i</b>	91	125
10	4-MeOC <sub>6</sub> H <sub>4</sub>	C <sub>2</sub> H <sub>5</sub>	<b>2j</b>	92	94
11	4-MeOC <sub>6</sub> H <sub>4</sub>	C <sub>3</sub> H <sub>7</sub>	<b>2k</b>	92	122
12	4-ClC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	<b>2l</b>	90	92

<sup>a</sup> Reaction conditions: *N,S*-acetal (5.0 mmol), allyl bromide (5.0 mmol), CuCl<sub>2</sub> (5.0 mmol), THF (30 mL), 3 h, reflux.



Scheme 2.

tetrahydrofuran and generated *N,S*-acetals **3a–l** in very low yields (10–15%).<sup>13</sup>

During optimization of the reaction conditions, the effect of a number of factors was investigated, including the nature of the base, solvent, the temperature (ranging from 0 °C to reflux), substrate concentration and reaction time. The best yields were obtained using NaH/THF at 25 °C under N<sub>2</sub> and stirring for 10 h giving a 15% yield (Table 3).

Subsequently, *N,S*-acetals **3a–l** were submitted to possible intramolecular rearrangements in the presence of cupric chloride and Lewis acids (AlCl<sub>3</sub>, ZnCl<sub>2</sub>, EtO–BF<sub>3</sub>). However, under these conditions, only inseparable by-products were formed and none of the reactions yielded the β-keto enamides **2a–l** (Scheme 3). Thus the possible mechanism for the formation of the C-allylated products from N-allylated adducts **3a–l** via a 3-aza-Cope rearrangement was ruled out. This implied that C-allylation rather than N-allylation was highly favourable in presence of the metal halide and Lewis acid catalytic conditions.

Compounds **2** existed as enamides as evident from their spectral data.<sup>14</sup> Compounds **2** exhibited ethylenic proton

Download English Version:

<https://daneshyari.com/en/article/5281477>

Download Persian Version:

<https://daneshyari.com/article/5281477>

[Daneshyari.com](https://daneshyari.com)