

Molecular iodine: a powerful catalyst for the easy and efficient synthesis of quinoxalines

Shivaji V. More, M. N. V. Sastry, Chieh-Chieh Wang and Ching-Fa Yao*

Department of Chemistry, National Taiwan Normal University 88, Sec. 4, Tingchow Road, Taipei, Taiwan 116, ROC

Received 13 May 2005; revised 24 June 2005; accepted 8 July 2005

Abstract—Various biologically important quinoxaline derivatives were efficiently synthesized in excellent yields using inexpensive, nontoxic, and readily available bench top chemical, iodine in catalytic amount (10 mol%). Besides this, a systematic study was carried out to evaluate parameters such as solvent and catalyst loading. Several aromatic as well as aliphatic 1,2-diketones and aromatic 1,2-diamines, such as substituted phenylene diamines, tetra amines were further subjected to condensation using catalytic amounts of iodine to afford the products in excellent yield.

© 2005 Published by Elsevier Ltd.

1. Introduction

Among the various classes of nitrogen containing heterocyclic compounds, quinoxaline derivatives are important components of several pharmacologically active compounds.¹ Although rarely described in nature, synthetic quinoxaline ring is a part of a number of antibiotics such as echinomycin, leromycin, and actinomycin, which are known to inhibit the growth of Gram-positive bacteria and are also active against various transplantable tumors.² Besides this, it has been reported for their application in dyes,³ efficient electroluminescent materials,⁴ organic semiconductors,⁵ building blocks for the synthesis of anion receptor,⁶ cavitands,⁷ dehydroannulenes,⁸ and DNA cleaving agents.⁹ A number of synthetic strategies have been developed for the synthesis of substituted quinoxalines and the most common method is the condensation of an aryl 1,2-diamine with 1,2-dicarbonyl compounds in refluxing ethanol or acetic acid.¹⁰ However, many improved methods have been reported for the synthesis of quinoxalines using catalytic amounts of various metal precursors, acids, and zeolites.¹¹ In addition to the above catalytic methods, reports were also known with microwave¹² and solid phase synthesis.¹³ Nevertheless, most of these methods

suffer from unsatisfactory product yields, critical product isolation procedures, expensive and detrimental metal precursors, and harsh reaction conditions, which limit their use under the aspect of environmentally benign processes.


The use of molecular iodine in organic synthesis has been known for a long time. In recent years, molecular iodine has received considerable attention as an inexpensive, nontoxic, readily available catalyst for various organic transformations¹⁴ under mild and convenient conditions to afford the corresponding products in excellent yields with high selectivity. However, there is no example of quinoxaline synthesis using molecular iodine as a catalyst. As part of our on going interest, in the use of cheap and ecofriendly materials as catalysts for various organic transformations, we had the opportunity to look into the synthesis of quinoxalines using molecular iodine.

2. Results and discussion

In the beginning, a systematic study was carried out for the catalytic evaluation of iodine toward the synthesis of quinoxalines. Initially, a blank reaction was conducted using benzil and *o*-phenylenediamine in boiling ethanol, which resulted in the formation of a condensation product after 45 min (60% Y). With the same substrates, the reaction in ethanol, using catalytic amount of iodine at room temperature afforded the products in

Keywords: 1,2-Diketones; Quinoxalines; Iodine; Catalytic amount; Ecofriendly process.

* Corresponding author. Tel.: +886 229309092; fax: +886 229324249; e-mail: cheyao@cc.ntnu.edu.tw



Scheme 1.

quantitative yield during 15 min. Thus, in the absence of iodine, the reaction was slow and also requires refluxing conditions with unsatisfactory yields. We next investigated the effect of different solvents on the reaction rate as well as the yields of products. Almost all the solvents afforded the products in excellent yield with a variation in time period. Protic solvents like methanol and ethanol afforded the products in high yield within 15–20 min.

After screening different solvents, acetonitrile came out as the solvent of choice, which not only afforded the products in good yield, but also with higher reaction rates. In general, for polar solvents the reaction rate is high, whereas for nonpolar solvents it is little low (Scheme 1 and Table 1).

The method of iodine catalyzed synthesis of quinoxalines is very simple, efficient, clean, and without any other side products. We also evaluated the amount of iodine required for this transformation. As less as, 5 mol% of iodine can catalyze the reaction to the same

extent, but needs a little longer reaction times (>1 h). In the absence of iodine, the reaction takes more than 24 h to complete. In an optimized reaction condition, 1,2-diketone (1 mmol) and 1,2-diamine (1.2 mmol) in acetonitrile (0.5 mL) were mixed with iodine (0.10 mmol) and stirred at room temperature for 3–30 min. After the completion of the reaction (monitored by TLC), a simple work up affords the products in excellent yield¹⁵ (Scheme 2).

In order to evaluate the efficiency of this methodology, a number of 1,2-diketones and 1,2-diamines were further subjected to condensation using catalytic amount of iodine (Table 2).

With symmetrical aromatic diamines the reaction showed good product yields. Diamine, such as naphthalene-1,2-diamine, steric factors played a key role in affecting the rate of reaction and the reaction requires a longer time. With electron donating substituents in the amine part, increased yields of products were obtained, whereas the effect is reverse with electron withdrawing substituents. On the other hand, electron donating substituents associated with aromatic 1,2-diketone decreased the product yields and the effect is reverse with electron withdrawing groups. However, the variations in the yields were very little and both substituted aromatic diamines such as 4-chloro and 3-methyl gave the condensed products in excellent yields with different substituted 1,2-diketones. To check the versatility of this method, we have also subjected other than symmetrical 1,2-diketones, such as furil and 1-phenyl-1,2-propanedione, and obtained the products in excellent yields. In the case of unsymmetrical 1,2-diketones such as 1-phenyl-1,2-propanedione, different aromatic diamines delivered a 1:1 ratio of regioisomers in almost quantitative yield (Table 3, entries 7 and 8). In another variation, tetra amines such as 3,3',4,4'-tetraamino-1,1'-biphenyl underwent condensation with diketone (benzil) in the presence of catalytic amount of iodine (10 mol%) and afforded the product **5** in 85% yield¹⁶ (Scheme 3).

This methodology is very useful for the synthesis of a sterically hindered amine such as 2,3,2',3'-tetraphenyl-[6,6'] biquinoxaliny **5**.


Though the role of iodine is not clearly known, but it can act as a mild Lewis acid. A part from its acidity iodine plays a complex role in accelerating the dehydrative steps, and thus promotes the formation of products.¹⁷

Table 1. Effect of solvents for the iodine catalyzed synthesis of quinoxalines

Entry ^a	Solvent	Time	Yield ^b (%)
1	CH ₂ Cl ₂	15 min	95
2	MeOH	20 min	92
3	EtOH	15 min	90
4	Et ₂ O	30 min	91
5	THF	30 min	94
6	DMF	15 min	89
7	CH ₃ CN	2–3 min	98
8	EtOAc	30 min	85
9	PhCH ₃	3 h	93
10	PhH	4.5 h	91

^a All reactions were performed at 1 mmol scale using 10 mol% of iodine in 0.5 mL of solvent.

^b Column isolated yields.



Scheme 2.

Download English Version:

<https://daneshyari.com/en/article/5286979>

Download Persian Version:

<https://daneshyari.com/article/5286979>

[Daneshyari.com](https://daneshyari.com)