

Tuning the photoinduced electron transfer in near-infrared heptamethine cyanine dyes

Fengling Song, Xiaojun Peng,* Erhu Lu, Yanan Wang, Wei Zhou and Jiangli Fan

State Key Laboratory of Fine Chemicals, Dalian University of Technology, 158 Zhongshan Road, Dalian 116012, PR China

Received 24 November 2004; revised 18 April 2005; accepted 20 April 2005

Available online 31 May 2005


Abstract—An efficient photoinduced electron transfer (PET) system in near-infrared region was described. The PET in heptamethine cyanine dyes was tuned by changing the electron-donating ability of the substituent at the central position of the polymethine chain. 4-Aminophenylthio-substitution led to an efficient PET and the lowest fluorescence quantum yield. The acetylation, protonation or transition metal cation coordination of the amino group could recover fluorescence greatly via suppressing the PET. © 2005 Elsevier Ltd. All rights reserved.

Fluorescent sensors via the photoinduced electron transfer (PET) strategy have been of great interest due to their various prospective applications.¹ However, most of these PET systems work effectively in UV–visible region, few of them in near-infrared (NIR) region. An inspection of the Weller equation² regarding intramolecular PET processes clearly shows that it is difficult to find a PET system when the HOMO–LUMO energy gap is very small as the case of NIR fluorescent dyes.³ An additional problem for NIR sensing schemes is that the ultrafast radiationless deactivation of the excited state is very efficient in these dyes, which can be an obstacle to use them as sensors.⁴

Recently heptamethine cyanine dyes employed as fluorescence labels and sensors *in vivo*⁵ have attracted immense interest because their spectra reach the NIR

region where biological matrix exhibits the least absorption and auto-fluorescence background.⁶ Patonay and co-workers have developed heptamethine cyanine dyes with a rigid chloro cyclohexenyl ring in the methine chain (like dye **2** in Scheme 1), which can increase the photostability and fluorescence quantum yield.^{7,8} This structure also provides the dye with a reactive chloro-group for chemical substitution at the central position.⁹ By this substitution many heptamethine cyanine dyes as biosensors and fluorescent probes were obtained.^{6,7,10,11} But few of them were employed as PET sensors for the above mentioned reasons.

Here we report three heptamethine cyanine dyes with thio-substituents in the central position in which PET can be tuned by changing the electron-donating ability of the substituent. The thioether bond was used as the



Scheme 1. Synthesis of the heptamethine cyanine dyes.

Keywords: Heptamethine cyanine dyes; Fluorescence sensors; Near-infrared; Photoinduced electron transfer.

* Corresponding author. Tel.: +86 411 8899 3899; fax: +86 411 8899 3906; e-mail: pengxj@dlut.edu.cn

spacer, which was seldom found. The PET in dye **1b** could be inhibited when proton, zinc or iron ion was added to the system and the fluorescence was observed to recover.

The synthesis of dyes **1a–c** was outlined in Scheme 1 according to a similar method.^{12,13} Dye **2** was stirred with thiophenols in anhydrous DMF under argon at room temperature in the dark for 1 h. The green products **1a–c** were purified on C18-RP column using methanol–water mixture as the eluent and characterized by spectroscopic data.¹⁴

The maximum absorption and emission wavelengths (λ_{max}) of **1a–c** had a small red shift compared with their parent dye **2** (Table 1). Dye **1b** was found to have a lower fluorescence quantum yield ($\Phi_f = 0.0065$) than the others. But, after acetylation of the amino group the quenched fluorescence recovers completely (**1c**, $\Phi_f = 0.036$, close to **1a**, $\Phi_f = 0.038$).

When a high-energy amino group was introduced into the *para* position of the thiophenyl, the substituent moiety in **1b** had a sufficiently high HOMO energy level to transfer an electron to the HOMO of the excited fluorophore. The ΔG_{PET} value of **1b** calculated from its potentials according to Weller equation^{2,3} was found to be the smallest among all dyes. It means that **1b** has the strongest thermodynamic force to perform an efficient PET process. But after acetylation of the amino

group, the HOMO energy level of the moiety in **1c** decreased and was close to that in **1a**, so that PET process was inhibited and the fluorescence recovered. The weaker fluorescence of dye **1a** and **1c** than their parent dye **2** might also be attributed to the weak electron-donating capability of thiophenol group, which led to a weak PET process. In our previous work, the attachment of a carboxymethyl mercapto group, a electron-withdrawing group, in the position caused a stronger fluorescence ($\Phi_f = 0.096$).¹² The relationship between molecule structure and orbital energy level was concluded in Figure 1.

According to the mechanism of PET, the HOMO energy level of the amine will decrease after the amine being protonated or bound by cation.¹⁶ However, PET process can generally occur when the HOMO–LUMO gap of the fluorophore is relatively large like fluorescence sensors in UV–visible region. It is difficult to achieve effective PET as the HOMO–LUMO gap is small according to the Weller equation regarding PET processes, for the energy introduced into the NIR system by excitation is too small to reduce the fluorophore and oxidize the donor via intramolecular electron transfer. So, it seems efficient PET is less likely for NIR dyes.³

Fortunately, the fluorescence off-on switching based on PET mechanism had been observed in the new dyes. The fluorescence of dye **1b** in aqueous solution became stronger with the increase of HCl concentration (Fig. 2). The inverse trend was observed in the cases of dye **1a**, **1c** and **2**, for cyanine dyes can generally be decolorized by light and acid.¹⁷ These results indicated that the PET was inhibited after the protonation of the amine.

It is well known that heavy and transition metal cations possess intrinsic properties that they can usually quench the emission of organic lumophores.¹⁸ And cyanine dyes have been used as sensors for transition metal ions such as Fe(III), Co(II) and Cu(III) based on Förster type quenching.^{19,20} This kind of fluorescence quenching was also observed in the cases of dye **1a** and dye **1c** when zinc (II) was added. However, dye **1b** (Fig. 3) exhibited a fluorescence enhancement effect with the increase of concentration of ZnSO₄. Similar results were obtained when ZnSO₄ was displaced by FeCl₃, Al₂(SO₄)₃, Ce(NO₃)₃.

Table 1. Photophysical characteristics of dyes in water at 1×10^{-6} M

Dye	λ_{ab} (nm)	λ_{em} (nm)	ϵ ($\times 10^5$) ^a	Φ_f ^b	$E_{\text{ox}} - E_{\text{re}}$ (eV) ^c	ΔG_{PET} (eV)
1a	794	817	2.3	0.038	1.242	−0.380
1b	795	820	0.9	0.0065	1.212	−0.408
1c	798	823	1.6	0.036	1.225	−0.389
2	783	803	2.0	0.065	1.260	−0.384

^a Molar extinction coefficients are in $\text{cm}^{-1} \text{M}$ and in the maximum of the highest peak.

^b The fluorescence quantum yields were determined in reference to IR-125 in DMSO ($\Phi_f = 0.13$).¹⁵

^c E_{ox} and E_{re} , the oxidative and reductive, respectively, were detected by cyclic voltammetry.




Figure 1. Structure and molecule orbital diagrams of the heptamethine cyanine dyes.

Download English Version:

<https://daneshyari.com/en/article/5287921>

Download Persian Version:

<https://daneshyari.com/article/5287921>

[Daneshyari.com](https://daneshyari.com)