

PHOTOINDUCED OXYGENATION OF DEHYDROROTENONES (1)


Hiroshi Suginome, Tomonori Yonezawa and Tadashi Masamune

Department of Chemistry, Faculty of Science, Hokkaido University

Sapporo, Japan

(Received in Japan 8 August 1968; received in UK for publication 27 August 1968)

We report here the observation that the mild irradiation of the solution of dehydrorotenone (Ic) and its homologue in the presence of sodium borohydride at 25° readily affords rotenonone (IIc) and its homologue in moderate yields.



Thus, in a typical experiment, dehydrorotenone (Ic) (2) (50 mg) in dioxane-ethanol mixture (30 ml) containing sodium borohydride (13 mg) was irradiated by a 150 W high pressure mercury arc lamp as the light source for 24 hours. The only major product was identified as rotenonone (IIc) (2a,3). In sharp contrast with this, we find that irradiation in the absence of sodium borohydride produced

no isolable IIc. Best yield was obtained in the oxygenation of Ia in the presence of NaBH_4 . Under comparable conditions, rotenone or isorotenone was recovered unchanged. These results together with those on the homologous compounds are tabulated below.

TABLE I

Irradiated compounds	Yield (%) of 6H-rotoxin-6,12-diones or 6a,12a-dihydro-6H-rotoxin-6,12-diones formed (4)		
	a		b
Ia (2C,5)	IIa* (2C,5)	: 40	trace
Ib (2C)	IIb (2C)	: 9	trace
Ic	IIc	: 10	trace
Id (6)	IIId** (6)	: 16	trace
Rotenone (Ic:6a,12a-dihydro-)	IIc,6a,12a-dihydro-	: 0	0
Isorotenone (Ia:6a,12a-dihydro-)	IIa,6a,12a-dihydro-	: 0	0

a) In the presence of NaBH_4

b) In the absence of NaBH_4

* Irradiation for 44 hours

** Irradiation for 94 hours (benzene:dioxane 2:1)

In the autoxidation at a methylene group flanked either by an oxygen atom or an aromatic ring, the product is mostly secondary hydroperoxides (7) and a very few examples of the formation of ketones in sizable amounts have been recorded (8). The termination reaction of peroxy radicals with an α -hydrogen would seem complicated although evidence has been proposed for a certain case (9).

Although the role of borohydride in the present case is not clear, borohydride would reduce the concentration of free radical initiators and would prohibit some of the undesirable side reactions. It is certain that even in the presence of NaBH_4 the primary products of the present oxygenation would be secondary hydroperoxides at 6 position of 6H-rotoxin rings.

One of the probable roles of borohydride would be to destroy these hydroperoxides immediately after their formation to convert them into the hemiacetal

Download English Version:

<https://daneshyari.com/en/article/5342864>

Download Persian Version:

<https://daneshyari.com/article/5342864>

[Daneshyari.com](https://daneshyari.com)