

Accepted Manuscript

Title: Strain in strain-free benzenoid hydrocarbons: The case of phenanthrene

Author: Slavko Radenković Ivan Gutman Slađana Đorđević

PII: S0009-2614(15)00130-X
DOI: <http://dx.doi.org/doi:10.1016/j.cplett.2015.02.039>
Reference: CPLETT 32832



To appear in:

Received date: 22-1-2015
Revised date: 18-2-2015
Accepted date: 19-2-2015

Please cite this article as: Slavko Radenković, Ivan Gutman, Slađana Đorđević, Strain in strain-free benzenoid hydrocarbons: The case of phenanthrene, *Chemical Physics Letters* (2015), <http://dx.doi.org/10.1016/j.cplett.2015.02.039>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

- Strain energy of repulsion between the two bay H-atoms is examined.
- A method to estimate the energy of this strain was developed.
- Strain energy in phenanthrene was found to be about 7 kJ/mol.

Accepted Manuscript

Download English Version:

<https://daneshyari.com/en/article/5380068>

Download Persian Version:

<https://daneshyari.com/article/5380068>

[Daneshyari.com](https://daneshyari.com)