



The possible phase diagrams for the transverse Ising model with temperature-dependent parameters



Bayor Jude Simons^a, Baohua Teng^{a,*}, Shanyu Zhou^b, Liming Zhou^c, Xiangyu Chen^c, Minghe Wu^a, Hao Fu^a

^aSchool of Physical Electronics, University of Electronic Science and Technology of China, Chengdu 610054, PR China

^bSchool of Yingcai, University of Electronic Science and Technology of China, Chengdu 611731, PR China

^cSchool of Optoelectronic Information, University of Electronic Science and Technology of China, Chengdu 610054, PR China

ARTICLE INFO

Article history:

Received 10 May 2014

In final form 12 May 2014

Available online 16 May 2014

ABSTRACT

In this Letter the transition properties of the transverse Ising model with temperature-dependent parameters are investigated. By supposing the simple dependent relations of the interaction parameters on temperature, the phase diagrams are straightforwardly obtained, which may be used to describe the closed-loop behavior for the phase transition of the systems. In fact, the reentrant phase behavior of the system obtained by this Letter is to some extent coincident with the phenomena exhibited in some colloids and complex fluid mixtures as well as proteins.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

The study of complex fluids, like some colloids and fluid mixtures has been being an important topic in physics and chemistry, and the complexity and peculiarity of these systems have been widely investigated by the theoretical and experimental researchers [1–14]. Davies et al. studied the solubility property of nicotine-water system [6]. Walker et al. found that the phenomenon of reappearing phases is caused by the complex competition between entropy and energy in a solvent containing hydrogen bonds [7]. Moelbert et al. obtained the phase diagram for an aqueous solution of nonpolar solute particles by mean-field calculation [8]. These studies showed the complexity of the binary solutions, and indicated the strength of interaction parameters among the system may increase with temperature [9]. An increase of temperature will lead to the overcome of enthalpic contribution of the two different components [10]. Meanwhile some results generated the closed-loop phase diagrams of concentration against temperature. These shapes of phase diagrams implied that the systems may have more interesting and attractive phase behaviors. The earliest closed-loop phase diagram was obtained in the nicotine-water system [11]. Lang et al. studied the reentrant behavior of alkyl polyoxyethylene nonionic surfactants [12]. Nord et al. studied the critical phenomena in polyvinyl alcohol-acetate copolymer solutions [13]. Narayanan and Kumar reviewed some important developments regarding phenomenon of reentrant phase

transitions in binary and multicomponent liquid mixtures [14]. Recently Dias studied the cold denaturation of proteins within the framework of numerical simulations of the Mercedes-Benz model for water [15], and the system also exhibits the reappearing phase behavior. In fact, the phenomenon of reentrant phase transition is intrinsically novel and is discovered in amazingly diverse systems such as binary gases, liquid crystals, microemulsions, gels, granular superconductors, proteins, organometallic compounds, crystals [14–17].

Theoretically, Ising or Ising-like model has a long and venerable tradition in the applications to order–disorder systems, such as ferromagnetic or ferroelectric systems, binary alloys, lattice gases, complex fluid, and even proteins. In ferromagnetic or ferroelectric field, Ising model has a wide and solid application [18–22]. In proteins field, Poland and Scheraga, for example, discussed a comprehensive survey of these models to deal with the coil-to-helix transition in proteins [23]. Further Badasyan et al. studied the cooperativity and stability of a helix-coil transition using a similar Ising-like model [24]. Zamparo et al. show an example for the case of Ising-like model by studying the kinetics of the Wako–Saito–Munoz–Eaton model of protein folding [25]. In complex fluid field, Campi and Krivine applied usual Ising model with temperature-dependent effective interaction to complex fluids [26], and obtained some meaningful results which can mimic the shapes of the phase diagram of complex fluids. In this Letter we will study the reentrant phase behavior of the system by using the transverse Ising model (TIM). Considering the temperature dependences of two kinds of effective interaction parameters, we will produce

* Corresponding author.

E-mail address: phytbb@163.com (B. Teng).

possible phase diagrams with the reentrant phase behavior and density curves against temperature with closed-loop shape.

2. The model

The Hamiltonian of the transverse Ising model is [18–22].

$$H = -\sum_i \Omega S_i^x - \sum_{\langle ij \rangle} J S_i^z S_j^z \quad (1)$$

where S_i^x and S_i^z are the x - and z -components of a pseudospin-1/2 operator at site i in the lattice, and $\sum_{\langle ij \rangle}$ runs over only distinct nearest-neighboring pairs. Ω and J are interaction parameters, and usually called the transverse field and exchange parameters, respectively.

Within the framework of the mean-field theory, the z component of the pseudospin can be written as [18–22],

$$\langle S_i^z \rangle = (\sigma/2\omega_0) \tanh(\omega_0/2k_B T) \quad (2)$$

where $\omega_0^2 = \Omega^2 + \sigma^2$ and $\sigma = \sum_j J \langle S_j^z \rangle$. The ensemble average of the pseudo-spin $\langle S_i^z \rangle$ is understood as the order parameter of the system. In fact, $\langle S_i^z \rangle$ is used to describe the transition of the system from order ($\langle S_i^z \rangle \neq 0$) to disorder ($\langle S_i^z \rangle = 0$) state. For complex fluid system, we use usually the average occupation number or density ρ of the site i of the lattice to describe the transition properties [26], where $\rho = \langle S_i^z \rangle + 1/2$. In the following we can calculate the density ρ and analyze the transition behaviors for the square lattice system.

3. The numerical calculations and phase diagrams

Experimental studies in complex fluids demonstrated that the interaction parameters in the system may increase with increasing temperature [9,10]. Even in some crystal materials the interaction parameters showed positive relations with temperature [16,17]. Theoretically, by introducing the effective exchange interaction with temperature dependence to the usual Ising model, Campi and Krivine generated the phase diagrams of closed-loop shape, and described the reentrant phase behavior of the phase diagram of complex fluids [26]. Commonly, a microscopic Ising model cannot have a temperature-dependent parameter. However, introducing the effective model (here the T -dependence) to a microscopic theory will help to understand the physics of complicated systems in a simple way [26]. In complex fluid, for example, the solvent effect may be to bring about a renormalized, temperature-dependent effective parameter [27]. In the transverse Ising model, therefore, we suppose the effective exchange and effective transverse field parameters J and Ω have simple temperature-dependent relations as follows:

$$J = J_0 \left(\frac{T}{T_0} \right)^n \quad \text{and} \quad \Omega = \Omega_0 \left(\frac{T}{T_0} \right)^m \quad (3)$$

where T_0 are arbitrary constant. In the following the effective parameters J_0 , Ω_0 and $k_B T$ are reduced by $k_B T_0$, and notated still as J_0 , Ω_0 and t for simplicity. In this way, we can calculate the density and phase diagrams for the transverse Ising model with effective temperature-dependent parameters. The interest of these calculations lies in three aspects. First, it is convenient to obtain the temperature dependence of the density with the closed-loop shape under the framework of the mean-field approximation. Secondly, the four kinds of the possible phase transition diagrams describing the reentrant phase behavior of the system can be obtained in the same framework without introducing supplementary assumptions. Finally, not only the exchange but also transverse field parameters are considered as effective temperature-dependence parameters, therefore the phase transition properties have obvious differences with the previous work to a large extent.

Figure 1a shows the phase diagram of the effective exchange parameter J_0 against temperature t for different n . For n not larger than 1.0 (red and blue curves), when J_0 is very small, the system is at disordered state for any given temperature, but when J_0 is large, the system is ordered at lower temperatures and disordered at higher temperatures. This situation corresponds just to the usual phase diagram of the TIM with effective temperature-independent parameters. For the case of n larger than 1.0 (green curve), at higher temperatures the system is at disordered state for smaller values of J_0 and can be at ordered state for larger J_0 values. At lower temperatures, however, the system may be at disordered state for a large range of J_0 values. This is anomalous compared with the above property for the usual TIM with effective temperature-independent parameters because the usual system eventually ends at ordered state when J_0 is large enough. This anomaly could be attributed to the fact that at lower temperatures, the effective exchange parameter J increases with temperature so slowly that the contribution of effective exchange interaction to the ordered state is diminished to a large extent.

Meanwhile, the phase diagram shows a crossover point of temperature $t = 1$. As the parameter n increases the disordered region of the system increases at below this temperature and decreases at above this temperature.

In fact, the anomalous phase diagram corresponds to a closed-loop density curve against temperature, as shown in Figure 1b. The system is disordered at lower temperatures and becomes ordered at intermediate temperatures and again assumes disordered state at higher temperatures. As the effective exchange parameter J_0 decreases, the closed-loop region for the ordered state decreases. For the usual phase diagram, the corresponding density curves are shown in Figure 1c and d. The system is ordered at lower temperatures and disordered at higher temperatures. Maximum density of 1.0 or minimum one of 0.0 is obtainable when values of n are less than 1.0. Generally, as n decreases from Figure 1b–d (with constant J_0), on the one hand, the temperature region for the ordered state also decreases. On the other hand, the temperature region for the ordered state transforms from higher temperatures to lower temperatures, and their shape profiles change from closed-loop shape to inverted-U shape.

Figure 2a shows the phase diagram of n against temperature t for different J_0 . When J_0 is small (red curve), for small values of n , the system is at ordered state at lower temperatures and at disordered state at higher temperatures. For intermediate values of n , the system is always at disordered state for any given temperature. For larger values of n , the system is at disordered state at lower temperatures and ordered state at higher temperatures.

When J_0 is large (blue curve), for small and larger values of n , the system is consistent with the situation when J_0 is small (red curve) respectively. For intermediate values of n , however, the system is at disordered state at lower temperatures, ordered state at intermediate temperatures and disordered state at higher temperatures.

The corresponding density curves are shown in Figure 2b and c. For small values of n , the system is ordered at lower temperatures and disordered at higher temperatures and maximum density of 1.0 or minimum one of 0.0 is obtainable. For a large range of n greater than 1, the system is at a disordered state at lower temperatures for certain J_0 . This shows the same anomaly for the system as illustrated in Figure 1a. For intermediate values of n , the system is disordered at lower temperatures and becomes ordered at intermediate temperatures and again assumes disordered state at higher temperatures. Therefore, a closed-loop density curve was obtained. For a large value of n , the system is disordered at lower temperatures and becomes ordered at higher temperatures. Thus the figure changes from closed-loop shape to U shape.

Download English Version:

<https://daneshyari.com/en/article/5380795>

Download Persian Version:

<https://daneshyari.com/article/5380795>

[Daneshyari.com](https://daneshyari.com)