

NMR resonance splitting of urea in stretched hydrogels: Proton exchange and $^{1}H/^{2}H$ isotopologues

Philip W. Kuchel, Christoph Naumann, Bogdan E. Chapman, Dmitry Shishmarev, Pär Håkansson, George Bacskay, Noel S. Hush

PII: S1090-7807(14)00217-1
DOI: http://dx.doi.org/10.1016/j.jmr.2014.08.004
Reference: YJMRE 5488

To appear in: *Journal of Magnetic Resonance*

Received Date: 3 June 2014
Revised Date: 29 July 2014

Please cite this article as: P.W. Kuchel, C. Naumann, B.E. Chapman, D. Shishmarev, P. Håkansson, G. Bacskay, N.S. Hush, NMR resonance splitting of urea in stretched hydrogels: Proton exchange and $^{1}H/^{2}H$ isotopologues, *Journal of Magnetic Resonance* (2014), doi: http://dx.doi.org/10.1016/j.jmr.2014.08.004

NMR resonance splitting of urea in stretched hydrogels: proton exchange and $^{1}H/^{2}H$ isotopologues

Philip W. Kuchel[a], Christoph Naumann[a], Bogdan E. Chapman[a], Dmitry Shishmarev[a], Pär Håkansson[c], George Bacskay[b] and Noel S. Hush[a,b]

[a]*School of Molecular Bioscience, University of Sydney, NSW 2006, Australia*
[b]*School of Chemistry, University of Sydney, NSW 2006, Australia*
[c]*School of Chemistry, University of Southampton, SO17 1BJ, United Kingdom*

Running title: Urea in stretched gels

Correspondence: Philip W. Kuchel
School of Molecular Bioscience
Building G08
University of Sydney
NSW 2006
Australia
Phone: +61 2 9351 3709
Fax: +61 2 9351 4726
Email: philip.kuchel@sydney.edu.au

Keywords: ^{1}H NMR spectroscopy; ^{2}H NMR spectroscopy; [1,3-^{15}N]urea; chirality; gelatin gel; isotopomers; residual dipolar coupling; residual quadrupolar coupling; stretched gel

Abbreviations: NMR, nuclear magnetic resonance; RDC, residual dipolar coupling; RQC, residual quadrupolar coupling; nOe, nuclear Overhauser effect