

## Communication

Measurement of sample temperatures under magic-angle spinning from the chemical shift and spin-lattice relaxation rate of  $^{79}\text{Br}$  in KBr powder

Kent R. Thurber, Robert Tycko\*

Laboratory of Chemical Physics, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Building 5, Room 112, Bethesda, MD 20892-0520, USA

## ARTICLE INFO

## Article history:

Received 1 August 2008

Revised 22 September 2008

Available online 25 September 2008

## Keywords:

Solid state NMR

Magnetic resonance

Magic-angle spinning

Temperature calibration

## ABSTRACT

Accurate determination of sample temperatures in solid state nuclear magnetic resonance (NMR) with magic-angle spinning (MAS) can be problematic, particularly because frictional heating and heating by radio-frequency irradiation can make the internal sample temperature significantly different from the temperature outside the MAS rotor. This paper demonstrates the use of  $^{79}\text{Br}$  chemical shifts and spin-lattice relaxation rates in KBr powder as temperature-dependent parameters for the determination of internal sample temperatures. Advantages of this method include high signal-to-noise, proximity of the  $^{79}\text{Br}$  NMR frequency to that of  $^{13}\text{C}$ , applicability from 20 K to 320 K or higher, and simultaneity with adjustment of the MAS axis direction. We show that spin-lattice relaxation in KBr is driven by a quadrupolar mechanism. We demonstrate a simple approach to including KBr powder in hydrated samples, such as biological membrane samples, hydrated amyloid fibrils, and hydrated microcrystalline proteins, that allows direct assessment of the effects of frictional and radio-frequency heating under experimentally relevant conditions.

Published by Elsevier Inc.

## 1. Introduction

In nuclear magnetic resonance (NMR) with magic-angle spinning (MAS), the sample is generally at the center of a rapidly spinning rotor, preventing placement of a conventional temperature sensor at the sample position. The sample temperature can be strongly affected by frictional heating of the rotor (particularly at high MAS frequencies) or absorption of energy from radio-frequency (rf) electric fields (particularly in hydrated samples or other lossy samples). Additionally, when separate gas sources are used for MAS drive and bearings and for temperature control, the temperature of the gas that surrounds the MAS rotor may be uncertain. Temperature gradients across the MAS rotor and in the surrounding gas create further complications.

Several techniques for determination of internal sample temperatures have been proposed, including calibrations by known phase transition temperatures [1–5] and measurements of temperature-dependent chemical shifts for  $^{13}\text{C}$  in samarium acetate [6],  $^{15}\text{N}$  in TTAA [7],  $^1\text{H}$  in bulk water [8],  $^1\text{H}$  in methanol soaked into TTMSS [9],  $^{207}\text{Pb}$  in lead nitrate [10–14],  $^{119}\text{Sn}$  in  $\text{Sm}_2\text{Sn}_2\text{O}_7$  and other compounds [15,16],  $^{31}\text{P}$  in  $(\text{VO})_2\text{P}_2\text{O}_7$  [17], and  $^1\text{H}$  and  $^{31}\text{P}$  in aqueous solutions of TmDOTP [18]. Alternatively, the temperature dependence of a spin-lattice relaxation rate ( $1/T_1$ ) can be used, e.g., that of  $^7\text{Li}$  [19]. The shift of quartz crystal resonances [20] and

the  $^{35}\text{Cl}$  NQR frequency of  $\text{NaClO}_3$  [21,22] have also been proposed as useful temperature-dependent parameters.

In this Communication, we describe temperature measurements based on the chemical shift and  $T_1$  of  $^{79}\text{Br}$  in KBr powder. This method has a number of advantages, especially for  $^{13}\text{C}$  MAS NMR: (i) The  $^{79}\text{Br}$  NMR frequency differs from the  $^{13}\text{C}$  NMR frequency by only 0.4%, allowing the  $^{13}\text{C}$  channel of the NMR probe to be used with only minor retuning; (ii) The temperature range from 20 to 320 K or higher can be covered; (iii) The  $^{79}\text{Br}$  signal can be used for both temperature and magic-angle monitoring [23]; (iv) The  $^{79}\text{Br}$  signal is strong and  $T_1$  is short at temperatures above 40 K. Small quantities of KBr are therefore sufficient, which can be included in the sample without sacrificing much sample volume as described below. Additionally, KBr is not highly toxic.


## 2. Temperature-dependent spin-lattice relaxation

Fig. 1a shows the temperature dependence of  $T_1$  for  $^{79}\text{Br}$  in KBr.  $T_1$  changes by almost three orders of magnitude from 100 to 15 K, providing very good temperature sensitivity. Although the temperature dependence is weaker at higher temperatures, the fast spin-lattice relaxation ( $T_1 = 75 \pm 3$  ms at 296 K) allows rapid measurement with high signal-to-noise, so that  $T_1$  measurements can be used to determine sample temperatures up to at least 320 K.

Fig. 1b compares  $T_1$  values for  $^{79}\text{Br}$  and  $^{81}\text{Br}$ . Both are spin-3/2 isotopes with high abundance and similar gyromagnetic ratios, allowing comparative measurements without modification of the

\* Corresponding author. Fax: +1 301 496 0825.

E-mail address: [robertty@mail.nih.gov](mailto:robertty@mail.nih.gov) (R. Tycko).




**Fig. 1.** (a)  $^{79}\text{Br}$  spin-lattice relaxation time  $T_1$  in KBr powder (static sample) as a function of temperature. The solid line is a theoretical expression for quadrupolar relaxation due to phonons [27,28], with the overall normalization fit to the experimental result at 70 K. The dashed line is the empirical expression given in the text. (b) Ratio of  $T_1$  values for  $^{79}\text{Br}$  and  $^{81}\text{Br}$  in KBr as a function of temperature. Solid and dashed lines show the expected ratios for quadrupolar and magnetic relaxation, respectively.  $T_1$  values were measured at 9.4 T.

NMR probe. For quadrupolar relaxation,  $1/T_1$  is proportional to the quadrupole interaction squared, *i.e.*,  $1/T_1 \propto \nu_Q^2$ ; for magnetic relaxation,  $1/T_1$  is proportional to the magnetic interaction squared, *i.e.*,  $1/T_1 \propto \gamma^2$  [24]. Thus, the ratio of the  $^{81}\text{Br}$  relaxation rate to the  $^{79}\text{Br}$  relaxation rate is predicted to be 0.701 for quadrupolar relaxation and 1.16 for magnetic relaxation [25]. Data in Fig. 1b demonstrate that spin-lattice relaxation in KBr is dominated by quadrupolar interactions down to at least 20 K. This result suggests that paramagnetic impurities, which might vary between samples, do not play a significant role. In fact, our  $T_1$  measurements on KBr powder ground with a mortar and pestle are consistent with previous measurements on an optically pure single crystal [26].

The experimental  $T_1$  data for  $^{79}\text{Br}$  fit the following empirical expression (dashed line in Fig. 1a) to within 5% between 20 K and 296 K:  $T_1 = 0.0145 + 5330T^{-2} + (1.42 \times 10^7)T^{-4} + (2.48 \times 10^9)T^{-6}$  (seconds). This empirical expression is similar to the theoretical temperature dependence for quadrupolar relaxation using a Debye distribution for lattice phonons. The solid line in Fig. 1a shows the theoretical expression with only one adjustable parameter, the overall normalization of the relaxation rate [27,28]. Well above the Debye temperature of  $\Theta_D = 177$  K [26,29],  $T_1$  is proportional to  $T^{-2}$  in theory, consistent with the experimental results. Below the Debye temperature, phonons are frozen out, and the temperature dependence is stronger. In the low-temperature limit,  $T_1$  is proportional to  $T^{-7}$  in theory [28]. However, this limit is only a good approximation at temperatures below our measurement range ( $T < 0.02 \Theta_D = 3.5$  K) [28].

$T_1$  data in Fig. 1 were measured at 9.4 T in a non-spinning helium-cooled probe, based on a customized Janis SuperTran continuous flow cryostat, that provides accurate sample temperatures [30–32] using both Cernox and platinum resistor temperature sensors (Lake Shore Cryotronics). The  $T_1$  values were measured by saturation-recovery of the central NMR line (saturation with 5–80  $\pi/2$  pulses with 0.3–10 ms gaps), using a single-exponential fit. In general, spin-lattice relaxation of a nucleus with spin  $> 1/2$  is not single-exponential, and the relaxation of the central and satellite transitions will be different [33]. However, for KBr under MAS, we have found the  $T_1$  values of the central NMR line and the MAS sidebands to be equal, both at room temperature and 25 K. We attribute this observation to fast spin exchange among the different transitions across the fairly narrow width ( $\sim 25$  kHz) of the quadrupolar-broadened lineshape. The spin-spin relaxation time of  $^{81}\text{Br}$  is 400  $\mu\text{s}$  at room temperature [34], and we have measured  $\sim 200$   $\mu\text{s}$  for  $^{79}\text{Br}$  at 30 K (non-spinning). Also,  $T_1$  for  $^{79}\text{Br}$  at different MAS frequencies  $\nu_{\text{MAS}}$  does not show any change other than that expected from frictional heating up to at least  $\nu_{\text{MAS}} = 16$  kHz (see Fig. 2).



**Fig. 2.** (a) Correlation of the  $^{79}\text{Br}$  chemical shift of KBr powder under MAS with that of  $^{207}\text{Pb}$  in lead nitrate, using identical conditions for the two measurements. The temperature scale is determined from the reported temperature dependence of the  $^{207}\text{Pb}$  chemical shift [10]. (b) Temperature dependence of the  $^{79}\text{Br}$  chemical shift of KBr powder under MAS, with the temperature scale determined by  $^{79}\text{Br}$   $T_1$  measurements. Inset shows the  $^{79}\text{Br}$  chemical shift in a static sample at lower temperatures. Solid circles in (a) and (b) are data in which heated or cooled nitrogen gas was used to vary the sample temperature. Open circles are data in which the sample temperature was raised above room temperature by MAS at various spinning frequencies. Solid lines are linear fits. Chemical shifts are normalized to 0.0 ppm at 296 K (100.025761 MHz for  $^{79}\text{Br}$ , 83.227189 MHz for  $^{207}\text{Pb}$ ).

Download English Version:

<https://daneshyari.com/en/article/5406967>

Download Persian Version:

<https://daneshyari.com/article/5406967>

[Daneshyari.com](https://daneshyari.com)