



Contents lists available at SciVerse ScienceDirect

Journal of Molecular Liquids

journal homepage: [www.elsevier.com/locate/molliq](http://www.elsevier.com/locate/molliq)

## Light scattering study of human serum albumin in pre-denaturation: Relation to dynamic transition in water at 42 °C

Vitaliy Bardik <sup>a,\*</sup>, Vladimir Gotsulskii <sup>b</sup>, Evgen Pavlov <sup>c</sup>, Nikolai Malomuzh <sup>b</sup>, Dmitry Nerukh <sup>c</sup>, Igor Yanchuk <sup>d</sup>, Serhiy Lavoryk <sup>d</sup>

<sup>a</sup> Kiev National Taras Shevchenko University, Physics Faculty, Prospect Acad. Glushkova, 4, Kiev 03127, Ukraine

<sup>b</sup> Odessa Mechnikov National University, Dvoryanskaya str., 2, Odessa 65026, Ukraine

<sup>c</sup> Aston University, Non-linearity and Complexity Research Group, Birmingham, B4 7ET, UK

<sup>d</sup> Nanomedtech, Gor'kogo str., 68, Kiev 03150, Ukraine

### ARTICLE INFO

#### Article history:

Received 24 February 2012

Received in revised form 6 September 2012

Accepted 14 September 2012

Available online xxxx

#### Keywords:

Dynamic light scattering

Conformational dynamics

Human serum albumin

Hydrogen bonds

Hydrodynamic radius

### ABSTRACT

Protein functional motions are ultimately connected to water dynamics. The goal of this study is to link the conformational dynamics of albumin to a dynamic transition taking place at ~42 °C in water. We report the results of dynamic light scattering measurements of albumin aqueous solution in the temperature interval 20–65 °C. The processing of the experimental data produced the temperature dependence of the macromolecular hydrodynamic radius. We demonstrate that the growth of the macromolecular size in this temperature range can be divided into two stages that are connected to the dynamical properties of water.

© 2012 Elsevier B.V. All rights reserved.

### 1. Introduction and background

Protein conformation and its dynamics define protein's biological functioning. Despite very intensive research the molecular mechanisms and the driving forces of conformational changes are largely unknown. What is generally accepted is the key role of water in protein conformational dynamics.

Frauenfelder and colleagues have experimentally shown that protein dominant conformational motions are slaved by the hydration shell and the bulk solvent [1], while the protein molecule itself provides an 'active matrix' necessary for guiding the water dynamics towards the biologically relevant conformational changes (the protein's 'function' or folding). The change in water dynamics at the shell of up to 20 Å (almost a dozen water molecule diameters) around proteins has been demonstrated in the work [2]. Many other experimental and simulation works support the fact of ultimate connection between liquid water dynamical properties and protein unique conformational motions [3–6]. The specific property of water and other glass-forming liquids responsible for this behavior is their 'dynamic heterogeneity' manifested in the long known, but still poorly understood  $\alpha$ - and  $\beta$ -relaxation phenomena in liquid and supercooled water [7,8]. The essence of the phenomenon is the existence of spatially separated areas in the liquid with qualitatively different dynamical properties dictated by

collective intermolecular interactions and in the case of water by the peculiarities of the hydrogen bonding network.

In comparison with the majority of one component liquids water shows many unusual properties which were the subject of extensive studies for many years [9,10]. It was shown in [11], by separating the contributions of different physical nature, that the behavior of the normalized shear viscosities for water and argon can be compared using the corresponding states principle. The analysis of the temperature dependences of the reduced shear viscosity of water and argon has revealed the temperature point  $T_H = 42$  °C, from which the behavior of these substances becomes similar (Fig. 1). The kinematic shear viscosity of liquids is one of the main transport coefficients. It is formed by different components of the thermal motion of molecules, in the first place, by the translational and rotational degrees of freedom. For water the significant influence on the manifestation of its properties is produced by hydrogen (H) bonds. Thus, if a molecule is connected with its nearest neighbors by three or four H-bonds, it can only oscillate around some temporary equilibrium position.

Analogous results have been obtained on the basis of the temperature dependencies analysis of the normalized fraction of volume and the normalized heat of evaporation for water and argon (Figs. 2 and 3) [12].

For the molecular interpretation of these data it was proposed that the molecular motion below  $T_H$  has a crystal like character. The descriptive criterion for this region is the validity of the strong inequality  $\tilde{\tau} = \frac{\tau_0}{\tau_1} \gg 1$ , where  $\tau_0$  is the residence time or the time of small

\* Corresponding author.

E-mail address: [vital@univ.kiev.ua](mailto:vital@univ.kiev.ua) (V. Bardik).




Fig. 1. Temperature dependence of the reduced viscosity of water (1) and argon (2), temperature is in reduced units,  $T_H \approx 0.55$  (taken from [11]).

oscillations of a molecule near its temporary equilibrium position [13],  $\tau_1$  is the time of transition to another temporary equilibrium position:  $\tau_1 \approx \frac{1}{\sqrt{1/kT}} \approx 5 \cdot 10^{-13}$  s. This situation is illustrated schematically in Fig. 4. Thus, the derivation of the temperature dependence of  $\tilde{\tau}$  allows to establish the region of the crystal like representation of water. The detailed analysis of  $\tilde{\tau}$  as a temperature function has been carried out on the basis of the quasi-elastic incoherent neutron scattering data in [14,15].

In the vicinity of  $T_H$  the ratio of the characteristic times tends to 1. We, therefore, suggest that the character of the thermal motion in water at  $T_H = (315 \pm 3)$  K undergoes a especial transformation, which will be qualified by us as a dynamic transition. The rapid growth of the residence time  $\tau_0$  with the temperature decay can be naturally interpreted

in the framework of cluster representations, especially characteristic for supercooled states. Indeed, it had been shown in [16] that the relative volume occupied by the crystal like clusters increases from the value  $\varphi = 0.11$  at the melting temperature  $T_m = 273$  K up to  $\varphi = 0.41$  at  $T = 243$  K. For these states of water  $\tau_0$  can be identified as the lifetime of the crystal like clusters. Their average size changes more slowly and remains close to 10 Å. Note that among clusters, the leading role belongs to the hexagonal rings, which are the building elements for the ordinary (hexagonal) ice. Probably,  $\varphi \rightarrow 0$  when  $T \rightarrow T_H$ . The crystal like picture of the thermal motion in water near the melting point is also supported by the results of computer simulations presented in [17]. There it was shown that for  $T < 284$  K the increment of the mean square displacement of a molecule is close to zero in the time interval  $10^{-13} \text{ s} < \tau_0$ , where  $\tau_0 > 10^{-12}$  s starting from  $T \leq T_m$ .

These data testify that the global hydrogen bonding network disintegrates into an ensemble of weakly interacting clusters: dimers, trimers, tetramers, etc. [15]. In other words, the spatial connectivity between linear molecular chains is disrupted in the vicinity of  $T \approx T_H$ . All these facts lead to the conclusion that the temperature of 42 °C is the point of a *dynamic transition* in water.

The value of  $T_H$  coincides with the physiological temperature at which the thermal denaturation of proteins takes place. As denaturation (unfolding) is a form of conformational dynamics of proteins, the dynamic transition can have an effect on the protein's elementary conformational changes.

The goal of the article is to relate the conformational dynamics of proteins at the level of macromolecular size to the dynamic properties of water (within the framework of the 42 °C hypothesis describing the transition when the 'liquid like' molecular motion changes to the 'gas like' one) in the temperature range 25 °C–60 °C, especially in the vicinity of  $T_H \approx 42$  °C as the threshold temperature of protein conformational stability.

Here we report the results of studying aqueous human serum albumin solution (HSA). HSA is a single chain protein with 585 amino acids, with a molecular weight of ~67,000 Da. The structure of this protein has been determined by X-ray crystallography of high resolution [18]. HSA plays a special role in transporting metabolites and drugs throughout the vascular system and also in maintaining the pH and osmotic pressure of plasma [19]. HSA structure and dynamics are known to be influenced by several factors, like pH, temperature, and binding of different ligands [20]. There is a number of investigations of HSA by fluorescent [21], IR [22], and NMR spectroscopies [23]. However, to the best of our knowledge, there are no studies of the connection between the size and structure of the protein molecule and its hydration, especially at different temperatures.

## 2. Methodology, results, and discussion

Aqueous solution of HSA at concentrations 1 wt.% with ions  $K^+$  and  $Na^+$  5 mmol/l was investigated by dynamic light scattering technique (DLS), also known as photon correlation spectroscopy (PCS) [24]. This technique has been successfully applied in studies of growth of fractal aggregates in water solutions of albumin and DNA [25], aqueous solutions of poly(ethylene glycol) [26], diffusive processes of alcohols and semidilute polymer solutions [27–29], cluster growth in aqueous sugars and glass-forming aqueous glucose solutions [30,31]. The measurements have been performed in the temperature interval 25 °C–50 °C with the 1 °C step. The vertically polarized light 632.8 nm from a helium-neon laser was focussed onto the sample (the beam diameter in the scattering volume 0.3 mm, scattering is regulated by the vertical slit diaphragm) and the light scattered at 45° was recorded by a photomultiplier. The diaphragm restricting the aperture of a scattering beam was located before photocathode. The homodyne detection was used in our study. The photoimpacts were digitized and fed to a correlator which computed intensity–intensity correlation function, which, for monodisperse samples consisting of a single particle size




Fig. 2. Temperature dependencies of the normalized fraction volume for water and argon: a)  $\tilde{\nu}(t) = v(t)/v_c$ , b)  $\tilde{\nu}(t) = v(t)/v_R$ . The light line in a) corresponds to  $0.86 \cdot \tilde{\nu}_R^{(Ar)}(t)$  (taken from [12]).

Download English Version:

<https://daneshyari.com/en/article/5412259>

Download Persian Version:

<https://daneshyari.com/article/5412259>

[Daneshyari.com](https://daneshyari.com)