

Author's Accepted Manuscript

Mars Topography Investigated Through the Wavelet Leaders Method: a Multidimensional Study of its Fractal Structure

Adrien Delière, Thomas Kleyntssens, Samuel Nicolay



PII: S0032-0633(16)30003-4
DOI: <http://dx.doi.org/10.1016/j.pss.2016.12.008>
Reference: PSS4259

To appear in: *Planetary and Space Science*

Received date: 6 January 2016
Revised date: 11 October 2016
Accepted date: 21 December 2016

Cite this article as: Adrien Delière, Thomas Kleyntssens and Samuel Nicolay Mars Topography Investigated Through the Wavelet Leaders Method: a Multidimensional Study of its Fractal Structure, *Planetary and Space Science* <http://dx.doi.org/10.1016/j.pss.2016.12.008>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and a review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

Mars Topography Investigated Through the Wavelet Leaders Method: a Multidimensional Study of its Fractal Structure

Adrien Delière^{a,1,*}, Thomas Kleyntssens^{a,1}, Samuel Nicolay^a

^a*Institute of Mathematics, University of Liège, Belgium*

Abstract

This work examines the scaling properties of Mars topography through a wavelet-based formalism. We conduct exhaustive one-dimensional (both longitudinal and latitudinal) and two-dimensional studies based on Mars Orbiter Laser Altimeter (MOLA) data using the multifractal formalism called Wavelet Leaders Method (WLM). This approach shows that a scale break occurs at approximately 15 km, giving two scaling regimes in both 1D and 2D cases. At small scales, these topographic profiles mostly display a monofractal behavior while a switch to multifractality is observed in several areas at larger scales. The scaling exponents extracted from this framework tend to be greater at small scales. In the 1D context, these observations are in agreement with previous works and thus suggest that the WLM is well-suited for examining scaling properties of topographic fields. Moreover, the 2D analysis is the first such complete study to our knowledge. It gives both a local and global insight on the scaling regimes of the surface of Mars and allows to exhibit the link between the scaling exponents and several famous features

*adrien.deliege@ulg.ac.be

¹The first two authors contributed equally to this work and are listed alphabetically.

Download English Version:

<https://daneshyari.com/en/article/5488049>

Download Persian Version:

<https://daneshyari.com/article/5488049>

[Daneshyari.com](https://daneshyari.com)