


## Accepted Manuscript

Transverse relaxation of cerebrospinal fluid depends on glucose concentration

A. Daoust, S. Dodd, G. Nair, N. Bouraoud, S. Jacobson, S. Walbridge, D.S. Reich, A. Koretsky



PII: S0730-725X(17)30153-4

DOI: doi: [10.1016/j.mri.2017.08.001](https://doi.org/10.1016/j.mri.2017.08.001)

Reference: MRI 8815

To appear in:

Received date: 18 April 2017

Revised date: 28 July 2017

Accepted date: 2 August 2017

Please cite this article as: A. Daoust, S. Dodd, G. Nair, N. Bouraoud, S. Jacobson, S. Walbridge, D.S. Reich, A. Koretsky, Transverse relaxation of cerebrospinal fluid depends on glucose concentration, (2017), doi: [10.1016/j.mri.2017.08.001](https://doi.org/10.1016/j.mri.2017.08.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Title**

Transverse relaxation of cerebrospinal fluid depends on glucose concentration

**Author names and affiliations**

A. Daoust<sup>1</sup>, S. Dodd<sup>1</sup>, G. Nair<sup>1</sup>, N. Bouraoud<sup>1</sup>, S. Jacobson<sup>1</sup>, S. Walbridge<sup>1</sup>, D. S. Reich<sup>1</sup> and A. Koretsky<sup>1</sup>

<sup>1</sup> National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD 20892, United States

Alexia Daoust, alexia.daoust@nih.gov  
 Steven Dodd, doddst@ninds.nih.gov  
 Govind Nair, govind.bhagavatheeshwaran@nih.gov  
 Nadia Bouraoud, bouraoudn@ninds.nih.gov  
 Stephen Jacobson, JacobsonS@ninds.nih.gov  
 Stuart Walbridge, stuartwalbridge@mail.nih.gov  
 Daniel S. Reich, reichds@ninds.nih.gov  
 Alan Koretsky, KoretskyA@ninds.nih.gov

**Corresponding author**

Dr Alexia Daoust, PhD  
 Institute of Neurological Disorders and Stroke,  
 National Institutes of Health,  
 Building 10, room ID52,  
 10 center drive,  
 Bethesda, MD 20892, United States  
 Tel: 301-594-7314  
 Email: [alexia.daoust@nih.gov](mailto:alexia.daoust@nih.gov)

**Abstract words count: 244**

**Text words count: 5410**

**Tables: 2**

**Figures: 6**

**Supplemental information: 4**

**References: 40**

**Key-words**

MRI, relaxation time, CSF, human, monkey, rodent

**Abbreviations**

CNR: contrast to noise ratio  
 CPMG: multiecho Car-Purcell-Meiboom-Gill  
 CSF: cerebrospinal fluid  
 T<sub>1</sub>: longitudinal relaxation time  
 T<sub>2</sub>: transversal relaxation time

**ACKNOWLEDGEMENTS**

This study was supported by the Intramural Research Program of the National Institute of Neurological Disorders and Stroke (NINDS), National Institutes of Health.

Download English Version:

<https://daneshyari.com/en/article/5491430>

Download Persian Version:

<https://daneshyari.com/article/5491430>

[Daneshyari.com](https://daneshyari.com)