

Jet evolution in a dense medium: event-by-event fluctuations and multi-particle correlations

Miguel A. Escobedo^{a,b}, Edmond Iancu^a

^a*Institut de Physique Théorique, Université Paris Saclay, CNRS, CEA, F-91191, Gif-sur-Yvette, France*

^b*Department of Physics, P.O. Box 35, 40014 University of Jyväskylä, Finland*

Abstract

We study the gluon distribution produced via successive medium-induced branchings by an energetic jet propagating through a weakly-coupled quark-gluon plasma. We show that under suitable approximations, the jet evolution is a Markovian stochastic process, which is exactly solvable. For this process, we construct exact analytic solutions for all the n -point correlation functions describing the gluon distribution in the space of energy [1, 2]. Using these results, we study the event-by-event distribution of the energy lost by the jet at large angles and of the multiplicities of the soft particles which carry this energy. We find that the event-by-event fluctuations are huge: the standard deviation in the energy loss is parametrically as large as its mean value [1]. This has important consequences for the phenomenology of di-jet asymmetry in Pb+Pb collisions at the LHC: it implies that the fluctuations in the branching process can contribute to the measured asymmetry on an equal footing with the geometry of the di-jet event (i.e. as the difference between the in-medium path lengths of the two jets). We compute the higher moments of the multiplicity distribution and identify a remarkable regularity known as Koba-Nielsen-Olesen (KNO) scaling [2]. These predictions could be tested via event-by-event measurements of the di-jet asymmetry.

1. Introduction

One of the observables in which the formation of a collective medium in heavy-ion collisions manifests in a very clear way is the dijet asymmetry, the energy difference between two approximately back-to-back jets [3, 4]. The usual interpretation of this observation is the following:

- The two energetic jets are initially created in a hard process, due to momentum conservation the two jets will have back-to-back directions and approximately the same energy.
- For simplicity we consider central collisions in which the interaction plane has rotational symmetry. The point of the collision region in which the hard process takes place is not always the center, a deviation from this point will have as a consequence that the effective size of the medium seen by each jet will be different.

Email address: miguel.a.escobedo-espinoza@jyu.fi (Miguel A. Escobedo)

- The two jets will lose energy inside of the medium, however the amount of energy loss will depend on the size of the medium that they traverse.

However, this might not be the whole story. In the previous discussion we were assuming that the energy loss is always the same at fixed medium size, in other words, we were neglecting fluctuations. The question is then, how big are these fluctuations? This is one of the main problems we are going to address in this proceedings and the answer we are going to find is that the typical deviation in the energy loss is of the order of the average value and therefore fluctuations can not be neglected. We are going to arrive to this conclusion by performing an analytical computation based on the results obtained in [5, 6]. A similar result was obtained recently by a Monte Carlo computation in [7]. Another issue we want to discuss in this proceedings is what are the event-by-event properties of gluons produced by the energy loss mechanism.

2. Jet quenching formalism

We are going to perform the computation using the BDMPS-Z theory [8, 9]. We make the approximation that parton virtualities are zero, therefore there would be no branching in absence medium. In this formalism all the information that we need from the medium is encoded in its length L and a parameter called $\hat{q}$. There are two time-scales that have a very important role in this problem. The first one we are going to discuss is the formation time. This is given by the uncertainty relation $\tau_f \sim (2\omega)/k_\perp^2$ where ω is the energy of the gluon that is being emitted. In a medium $k_\perp \sim \sqrt{\hat{q}\tau_f}$, this gives a self consistent equation that results in $\tau_f \sim \sqrt{(2\omega)/\hat{q}}$. Another important time-scale is the branching time. In the BDMPS-Z theory the probability to emit a gluon during a small time Δt is

$$P(\omega, \Delta t) \propto \frac{N_c \alpha_s}{\pi} \sqrt{\frac{\hat{q}}{\omega}} \Delta t, \quad (1)$$

the branching time τ_{br} is the period after which we are almost sure that a gluon with a given energy will be emitted, looking at the previous equation we can see that $\tau_{br}(\omega) = \frac{\pi}{N_c \alpha_s} \tau_f(\omega)$, this shows that in perturbation theory the formation time is much smaller than the branching time and therefore, at first approximation, the branching process can be thought as an almost classical process in which gluons are formed instantaneously.

The branching time allows to divide the gluons in two different types. On one hand we have soft gluons that have an energy such that $\tau_{br} \ll L$ therefore they will be emitted abundantly. On the other hand we have the class of the hardest gluons which are likely to be emitted. They are those with $\tau_{br} \sim L$, this implies that they will have an energy of order $\omega_{br} \sim \alpha_s^2 \hat{q} L^2$. Their emission by the leading particle will dominate the energy loss. However, this gluons with energy ω_{br} will subsequently branch and at the end of the day what will be found is a lot of soft gluons emitted at large angles. The equations and the consequences of the multiple branching obtained with the previous assumptions were discussed in [5, 6], there it was observed the importance of the so-called democratic branching, the process in which a parton branches in a way such that the resulting partons have a similar energy. This will be a rare event for the leading particle because their energy is much bigger than ω_{br} , however for the gluons emitted by the leading particle, that will typically have an energy of the order of ω_{br} or smaller, this will be a very common process and a very efficient way to transfer energy into low energy gluons emitted at large angles.

3. The gluon spectrum and the average energy loss

The main focus of this section is going to be the gluon spectrum that we define as $D(x, t) = x \langle \sum_i \delta(x_i - x) \rangle$, where x is the energy fraction carried by the gluon. This quantity evolves with time following the equation [10]

$$\frac{\partial}{\partial \tau} D(x, \tau) = \int dz \mathcal{K}(z) \left[\sqrt{\frac{z}{x}} D\left(\frac{x}{z}, \tau\right) - \frac{z}{\sqrt{x}} D(x, \tau) \right], \quad (2)$$

where $\tau = \frac{\alpha_s N_c}{\pi} \sqrt{\frac{\hat{q}}{E}} t = \frac{t}{\tau_{br}(E)}$. E in this case is the energy of the leading particle. The case interesting for jet quenching at LHC is therefore the one in which $\tau \ll 1$, however the case $\tau \sim 1$ is also interesting in order

Download English Version:

<https://daneshyari.com/en/article/5493991>

Download Persian Version:

<https://daneshyari.com/article/5493991>

[Daneshyari.com](https://daneshyari.com)