



Higher harmonic anisotropic flow of identified particles in Pb–Pb collisions with the ALICE detector

Naghmeh Mohammadi for the ALICE Collaboration

Nikhef, Science Park 105, 1098 XG Amsterdam, The Netherlands

Abstract

Anisotropic flow plays a crucial role in establishing the equation of state of the Quark Gluon Plasma. The results at RHIC and LHC have demonstrated that the matter created in heavy-ion collisions behaves as a nearly perfect fluid reflected in the low value of the shear viscosity over entropy density ratio (η/s). The higher flow harmonics are particularly sensitive to the value of η/s in hydrodynamic calculations. In this analysis, we present the first ALICE results on the p_T differential v_2 , v_3 and v_4 for π^\pm , K^\pm , $p(\bar{p})$ from the high statistics 2011 heavy-ion run. We investigate how these v_n coefficients evolve with particle mass and centrality for the 0–1% and 20–30% centrality ranges. These new measurements aim at differentiating between models that use different initial conditions, constraining further the value of η/s and allowing to decouple the influence of the late hadronic stage from the hydrodynamic evolution of the system.

Keywords: RHIC, LHC, ALICE, anisotropic flow

1. Introduction

The main objective of heavy-ion collision experiments is to study the properties of the Quark Gluon Plasma [1]. Anisotropic flow, that characterises the momentum anisotropy of the final state particles, probes the properties of the system created in heavy-ion interactions, such as the equation of state and the ratio of shear viscosity to entropy density (η/s). This momentum anisotropy stems from the initial spatial anisotropy of the collision, reflected in its geometry, and by the initial density profile of nucleons participating in the collision which is different from one event to the other [2]. Anisotropic flow is characterized by a Fourier expansion of the azimuthal distribution of particle production relative to the symmetry plane [3], according to:

$$dN/d(\varphi - \Psi_n) \approx 1 + \sum_n 2v_n \cos[n(\varphi - \Psi_n)], \quad (1)$$

where φ is the azimuthal angle of particles, Ψ_n the azimuthal angle of the n th-order symmetry plane and v_n are the flow coefficients, with v_2 , v_3 and v_4 the elliptic, triangular and quadrangular flow, respectively. In this contribution, the results are reported for $v_2(p_T)$, $v_3(p_T)$ and $v_4(p_T)$ of identified charged pions, charged kaons and (anti-) protons in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV measured with the ALICE detector.

2. Analysis details

The data sample used for this analysis was recorded by ALICE during the 2011 Pb–Pb run at the LHC. Minimum-bias Pb–Pb events were triggered by the coincidence of signals from two forward detectors (V0A and V0C) positioned at both sides of the interaction point. In addition, an online selection based on the same V0 detectors was used to increase the number of central (i.e. 0–10% centrality range) and semi-central (i.e. 10–50% centrality range) events.

Events with a reconstructed primary vertex, whose position on the beam axis is within ± 10 cm from the nominal interaction point, were considered in this analysis. The data were grouped according to fractions of the hadronic cross section. The two centrality ranges presented in this article correspond to the 1% most central (i.e. smallest impact parameter) Pb–Pb collisions (~ 2.2 million) and the 20–30% interval (i.e. semi-central with large impact parameter) with 1.5 million events.

The charged particle reconstruction was done with the Inner Tracking System (ITS) and the Time Projection Chamber (TPC). The charged particle momenta were measured by the ITS and the TPC with a full azimuthal coverage in the pseudo-rapidity range $|\eta| < 0.9$. Charged pions and kaons as well as (anti-)protons were identified by combining the signals from the TPC and Time Of Flight (TOF) detectors similarly to the procedure followed in [4]. In this analysis, pions and protons are identified with a purity of $> 90\%$ in a transverse momentum range of $0.3 < p_T < 6$ GeV/c and kaons with a purity of $> 85\%$ in $0.3 < p_T < 4$ GeV/c. For more information about the ALICE detector and its performance see [5].

The flow harmonics were measured with the scalar product method introduced in [6]. A pseudo-rapidity gap of $|\Delta\eta| > 0$ was applied between the identified pions, kaons and (anti-)protons selected from one side of the TPC and the charged reference particles selected from the other side of the TPC. Due to this small pseudorapidity gap applied, correlations not related to the common symmetry plane, known as non-flow, contribute to the measured v_n^{AA} values. In this analysis, the contribution from non-flow was estimated using the HIJING event generator and is denoted as δ^{MC} [6]. The maximum magnitude of this correction is around 0.017 for v_2 , 0.008 for v_3 and 0.006 for v_4 in intermediate p_T region ($p_T > 3$ GeV/c). The reported values in this article are denoted as, v_n^{AA-MC} , and were extracted according to:

$$v_n^{AA-MC} = v_n^{AA} - \delta^{MC} \quad (2)$$

More details about the applied non-flow subtraction can be found in [6].

3. Results

Fig. 1 presents the p_T differential v_2^{AA-MC} , v_3^{AA-MC} and v_4^{AA-MC} for charged pions and kaons and the combination of protons and anti-protons, in the left, middle and right plot, respectively, for the 1% most central Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV (upper row). The bottom row of Fig. 1 presents the same picture that develops for the 20–30% centrality range. Note that the v_2 values for this centrality range were calculated with the scalar product method with $|\Delta\eta| > 0.9$ and without MC subtraction can be found in [4].

For the most central Pb–Pb collisions one expects the influence of the collision geometry to the development of v_n to be reduced as compared to the contribution of the initial density fluctuations. It is seen that for charged pions around 0.8 GeV/c, v_3 gets equal to v_2 and becomes the dominant harmonic beyond this value. Furthermore, at around 2 GeV/c, v_4 becomes equal to v_2 . For higher transverse momentum values, v_4 becomes gradually larger than v_2 reaching similar magnitude as v_3 at around 3.5 GeV/c. However, for the 20–30% centrality class where the collision geometry is expected to become the primary contributor to the development of v_n , it is seen that v_2 is the dominant harmonic throughout the entire measured momentum range. Furthermore, both v_3 and v_4 seem to have similar magnitudes and p_T evolution as the one observed for the most central Pb–Pb events, indicating a smaller influence of the collision geometry in their development than v_2 .

Download English Version:

<https://daneshyari.com/en/article/5494183>

Download Persian Version:

<https://daneshyari.com/article/5494183>

[Daneshyari.com](https://daneshyari.com)