



Contents lists available at ScienceDirect

Physics Letters A

www.elsevier.com/locate/pla



Dynamic dielectric and magnetic properties of a spontaneously polarized single layer of the polar dielectric

Vanik E. Mkrtchian^a, Hamlet G. Badalyan^b

^a Institute for Physical Research, Armenian Academy of Sciences, Ashtarak, 0203, Armenia

^b Department of General Physics, Yerevan State University, 1 Alex Manoogian St., Yerevan 0025, Armenia

ARTICLE INFO

Article history:

Received 7 August 2017

Received in revised form 9 September 2017

Accepted 11 September 2017

Available online xxxx

Communicated by Z. Siwy

Keywords:

Spontaneous polarization

Polar dipoles

Dielectric susceptibility tensor

Phospholipid bilayer

ABSTRACT

In the framework of the Debye–Smoluchowski theory and in local electric field approximation, an expression of the dielectric susceptibility tensor is derived for a single layer of spontaneously polarized polar dielectric in the field of a monochromatic radiation. It is shown that the AC radiation also generates or induces magnetic moments with a density that is expressed by the components of the dielectric susceptibility tensor of the layer.

© 2017 Published by Elsevier B.V.

1. Introduction

In equilibrium, the polar groups of phospholipid molecules, located on opposite sides of the cell membrane, are oriented mirror asymmetrically to each other [1–3]. By the method of minimization of phospholipid bilayers free energy, it was shown the existence in [1] of a finite inclination angle and lower and upper limits of its variation. Further, it was shown in [4] that the inclination angle of the dipoles strongly depends of the ratio of water to phospholipid concentrations.

The existence of spontaneously aligned rigid dipoles makes the system a strongly anisotropic biaxial material with electromagnetic properties completely different from the properties of the nematic liquid crystals [3]. In present theories of dielectric permittivity of a lipid bilayer [4], the contributions of the inclination of dipoles in the electromagnetic properties of cell membrane are ignored. This paper aims to investigate the problem of the influence of the spontaneous polarization on the dynamic, dielectric properties of the cell membrane in the field of a homogeneous AC radiation and so to eliminate the drawback in existing modern theories.

This paper is arranged as follows. It contains the theoretical part, which includes Sec. 2, of the mathematical description of the spontaneous polarization of a single layer of polar dipoles. In Sec. 3, we solve the Debye–Smoluchowski kinetic equation in the presence of an AC external field. The polarization tensor of a single




Fig. 1. Mirror asymmetric arrangement of the phospholipid molecules and single rigid dipole with inclination angle θ .

rigid dipole is found in Sec. 4, which is used in Sec. 5 to calculate the dielectric susceptibility in the local field approximation. In Sec. 6, we show a method of calculating the magnetic moment associated with the bounded dipole layer.

2. Spontaneous polarization

Because of the mirror symmetry arrangement of phospholipid molecules as seen in Fig. 1, we will consider single layers of spontaneously inclined dipoles.

The polarization of a rigid dipole is defined only by its direction \mathbf{n}

$$\mathbf{p} = p\mathbf{n}. \quad (1)$$

To handle mathematically the spontaneously alignment $\mathbf{p}^0 = p\mathbf{n}^0$ of the dipole parallel to the vector $\mathbf{n}^0 = (\sin\theta_0, 0, \cos\theta_0)$ in the

E-mail address: vem@daad-alumni.de (V.E. Mkrtchian).

<http://dx.doi.org/10.1016/j.physleta.2017.09.019>

0375-9601/© 2017 Published by Elsevier B.V.

plane xoz , we introduce a static, auxiliary electric field \mathbf{E}^0 acting in that direction

$$\mathbf{E}^0 = E^0 \mathbf{n}^0. \quad (2)$$

The dipole–field interaction U_0 is given by

$$U_0 = -\mathbf{p} \mathbf{E}^0. \quad (3)$$

The Boltzmann distribution of dipole moments in equilibrium at temperature T is

$$f^0 = C e^{-w_0(\mathbf{n}^0 \mathbf{n})}, \quad w_0 = pE^0/T, \quad (4)$$

where the coefficient C is defined from the normalization condition

$$\int f^0 d\mathbf{n} = 1 \quad (5a)$$

and it is expressed via a hyperbolic sine function

$$C = w_0 / (4\pi \sinh w_0). \quad (5b)$$

To assure the correctness of the description of the spontaneous polarization, we consider the limit $T \rightarrow 0$ of expression (4) for the distribution function f^0 . Using the limit representation of the Dirac function

$$\delta(x) = \frac{1}{2} \lim_{\varepsilon \rightarrow \infty} \varepsilon e^{-\varepsilon|x|}$$

we obtain

$$\lim_{T \rightarrow 0} f^0 = \frac{1}{\pi} \delta(1 - \mathbf{n}^0 \mathbf{n}), \quad (6)$$

which indicates that at temperature $T = 0$, the dipoles are “frozen” parallel to \mathbf{n}^0 .

In equilibrium at the finite temperature T , the dipole (1) fluctuates around, \mathbf{n}^0 . We can find, using expression (4) for distribution function, that the average polarization $\langle \mathbf{p} \rangle_0$

$$\langle \mathbf{p} \rangle_0 = \mathbf{p}^0 L(w_0), \quad (7a)$$

where L is the Langevin function

$$L(x) = \coth x - 1/x. \quad (7b)$$

Our result (7) is consistent with the mean electric moment calculated in [5].

3. The Debye–Smoluchowski equation

We first introduce the interaction V of the dipoles with a homogeneous and monochromatic electric field with the amplitude E and frequency ω

$$V = -\mathbf{E} d e^{i\omega t}. \quad (8)$$

Then total field–dipole interaction U is given by

$$U = U_0 + V. \quad (9)$$

The distribution function of rigid dipoles f obeys the Debye–Smoluchowski equation [5,6]

$$\partial_t f - D \Delta f - \zeta^{-1} [\nabla f \cdot \nabla U + f \Delta U] = 0, \quad (10)$$

where D , ζ are the diffusion and the friction coefficients, respectively.

In the absence of a time dependent electric field E , the system is in equilibrium and $D = T/\zeta$ and the distribution function f is given by f^0 in (4). In the presence of the field E , we describe

the distortions from the equilibrium by introducing a function F , defined by the product,

$$f = F f^0. \quad (11)$$

We restrict our consideration in domains of small external electric fields and then, in the linear approximation of V , substitution of (11) in the Debye–Smoluchowski equation (10) gives us the equation for F

$$\zeta \partial_t F + \nabla U_0 \cdot \nabla F - T \Delta F = \Delta V - T^{-1} \nabla U_0 \cdot \nabla V. \quad (12)$$

Expanding F in the basis of spherical functions

$$F = 1 + e^{i\omega t} \sum_{l,m} a_{l,m} Y_{l,m}(\mathbf{n}) \quad (13)$$

we obtain a set of nonhomogeneous algebraic equations for the amplitudes $a_{l,m}$. Because of the strong oscillating character of higher harmonics of $Y_{l,m}$, we may consider the contributions of only the first four harmonics $Y_{0,0}$ and $Y_{1,m}$ and so we obtain the following expression for F of (12)

$$F = 1 + \frac{p(\mathbf{n}E)}{T(1+i\omega\tau)} e^{i\omega t}. \quad (14)$$

In the absence of an initial polarization, that is, $w_0 = 0$, the response of the system to an external electric field is isotropic and that is equal to the corresponding Debye’s expression [5].

4. Polarization tensor

Consider the distribution function

$$f = f^0 + f^1. \quad (15)$$

For any quantity g , we introduce the notation $\langle g \rangle_{0,1}$ for the averages of g over f^0 and f^1 , respectively. We suppose f^0 to be normalized, i.e., $\langle 1 \rangle_0 = 1$ while the normalization integral of f^1 is $\langle 1 \rangle_1$, that is different of 1. In the linear approximation of f^1 , the average of g in the state (15) $\langle g \rangle$ is given by

$$\langle g \rangle = \langle g \rangle_1 - \langle g \rangle_0 \langle 1 \rangle_1. \quad (16)$$

Now, if we suppose the correction of f^1 to be proportional to f^0 (see (11) and (14)), then $f^1 = \lambda f^0$ and so

$$\langle g \rangle = \langle \lambda g \rangle_0 - \langle \lambda \rangle_0 \langle g \rangle_0 \quad (17)$$

for the average value of quantity g in linear approximation. In (17) $\langle \dots \rangle_0$ is the average in the 0-order distribution f^0 .

Defining the time dependent part of the dipole moment by

$$\langle \mathbf{p} \rangle(t) = \langle \mathbf{p}(\omega) \rangle e^{i\omega t},$$

where $\langle \mathbf{p}(\omega) \rangle$ is related to the polarization tensor α_{ij} by

$$\langle p_i(\omega) \rangle = \alpha_{ij}(\omega) E_j \quad (18)$$

and using (17) and (18), we find for the polarization tensor in the perturbed state (14) that

$$\alpha_{ij}(\omega) = \alpha(\omega) [\delta_{ij} + 2\beta n_i^0 n_j^0], \quad (19)$$

where

$$\alpha(\omega) = \frac{d^2 L}{w_0 T (1 + i\omega\tau)}, \quad (20)$$

$$2\beta = 3 + (w_0/L)(L^2 - 1) \quad (21)$$

and L is Langevin function (7b). In the limit $w_0 \rightarrow 0$, with $w_0^{-1} L \rightarrow 1/3$, we obtain

Download English Version:

<https://daneshyari.com/en/article/5496173>

Download Persian Version:

<https://daneshyari.com/article/5496173>

[Daneshyari.com](https://daneshyari.com)