

## Accepted Manuscript

On Wright's generalized Bessel kernel

Lun Zhang

PII: S0167-2789(16)30366-9

DOI: <http://dx.doi.org/10.1016/j.physd.2016.09.005>

Reference: PHYSD 31850

To appear in: *Physica D*

Received date: 19 July 2016

Accepted date: 20 September 2016



Please cite this article as: L. Zhang, On Wright's generalized Bessel kernel, *Physica D* (2016), <http://dx.doi.org/10.1016/j.physd.2016.09.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# On Wright's generalized Bessel kernel

Lun Zhang\*

September 26, 2016

## Abstract

In this paper, we consider the Wright's generalized Bessel kernel  $K^{(\alpha, \theta)}(x, y)$  defined by

$$\theta x^\alpha \int_0^1 J_{\frac{\alpha+1}{\theta}, \frac{1}{\theta}}(ux) J_{\alpha+1, \theta}((uy)^\theta) u^\alpha du, \quad \alpha > -1, \quad \theta > 0,$$

where

$$J_{a,b}(x) = \sum_{j=0}^{\infty} \frac{(-x)^j}{j! \Gamma(a + bj)}, \quad a \in \mathbb{C}, \quad b > -1,$$

is Wright's generalization of the Bessel function. This non-symmetric kernel, which generalizes the classical Bessel kernel (corresponding to  $\theta = 1$ ) in random matrix theory, is the hard edge scaling limit of the correlation kernel for certain Muttalib-Borodin ensembles. We show that, if  $\theta$  is rational, i.e.,  $\theta = \frac{m}{n}$  with  $m, n \in \mathbb{N}$ ,  $\gcd(m, n) = 1$ , and  $\alpha > m - 1 - \frac{m}{n}$ , the Wright's generalized Bessel kernel is integrable in the sense of Its-Izergin-Korepin-Slavnov. We then come to the Fredholm determinant of this kernel over the union of several scaled intervals, which can also be interpreted as the gap probability (the probability of finding no particles) on these intervals. The integrable structure allows us to obtain a system of coupled partial differential equations associated with the corresponding Fredholm determinant as well as a Hamiltonian interpretation. As a consequence, we are able to represent the gap probability over a single interval  $(0, s)$  in terms of a solution of a system of nonlinear ordinary differential equations.

## 1 Introduction and statement of results

### 1.1 Muttalib-Borodin ensembles and the Wright's generalized Bessel kernel

In 1995, Muttalib proposed the following class of joint probability density functions

$$\frac{1}{Z_n} \Delta(x_1, \dots, x_n) \Delta(x_1^\theta, \dots, x_n^\theta) \prod_{j=1}^n w(x_j), \quad \theta > 0, \quad x_j > 0, \quad (1.1)$$

for  $n$  particles  $x_1, \dots, x_n$ , where  $w$  is a weight function over the positive real axis,

$$Z_n = \int_{[0, \infty)^n} \Delta(x_1, \dots, x_n) \Delta(x_1^\theta, \dots, x_n^\theta) \prod_{j=1}^n w(x_j) dx_j$$

---

\*School of Mathematical Sciences and Shanghai Key Laboratory for Contemporary Applied Mathematics, Fudan University, Shanghai 200433, P. R. China. E-mail: lunzhang@fudan.edu.cn

Download English Version:

<https://daneshyari.com/en/article/5500358>

Download Persian Version:

<https://daneshyari.com/article/5500358>

[Daneshyari.com](https://daneshyari.com)