Walking speed decline in older adults is associated with elevated pro-BDNF in plasma extracellular vesicles

Caitlin Suire, Erez Eitan, Nancy Chiles Shaffer, Qu Tian, Stephanie Studenski, Mark P. Mattson, Dimitrios Kapogiannis

PII: S0531-5565(17)30347-9
DOI: doi: 10.1016/j.exger.2017.08.024
Reference: EXG 10131

To appear in: *Experimental Gerontology*

Received date: 10 May 2017
Revised date: 17 July 2017
Accepted date: 18 August 2017

Please cite this article as: Caitlin Suire, Erez Eitan, Nancy Chiles Shaffer, Qu Tian, Stephanie Studenski, Mark P. Mattson, Dimitrios Kapogiannis , Walking speed decline in older adults is associated with elevated pro-BDNF in plasma extracellular vesicles, *Experimental Gerontology* (2017), doi: 10.1016/j.exger.2017.08.024

Walking Speed Decline in Older Adults is Associated with Elevated Pro-BDNF in Plasma Extracellular Vesicles

Caitlin Suire[1*]; Erez Eitan[1*]; Nancy Chiles Shaffer[2*], Qu Tian[2], Stephanie Studenski[2], Mark P. Mattson[1] and Dimitrios Kapogiannis[1]

[1]Laboratory of Neurosciences, National Institute on Aging on Intramural Research Program, Baltimore, MD 21224. [2]Translational Gerontology Branch, National Institute on Aging on Intramural Research Program, Baltimore, MD 21224.

*These authors made approximately equal contributions to the generation and analyses of data.

Corresponding Author: D. Kapogiannis; 3001 S. Hanover St., NM531, Baltimore, MD, 21225; email: kapogiannisd@mail.nih.gov; tel: 410-350-3953

Acknowledgements: This work was supported entirely by the Intramural Research Program of the National Institute on Aging, NIH.