


UCP2 inhibition induces ROS/Akt/mTOR axis:
role of GAPDH nuclear translocation in
genipin/everolimus anticancer synergism

Ilaria Dando, Raffaella Pacchiana, Elisa Dalla
Pozza, Ivana Cataldo, Stefano Bruno, Paola Conti,
Marco Cordani, Anna Grimaldi, Giovanna Butera,
Michele Caraglia, Aldo Scarpa, Marta Palmieri,
Massimo Donadelli



www.elsevier.com

PII: S0891-5849(17)30775-X
DOI: <http://dx.doi.org/10.1016/j.freeradbiomed.2017.09.022>
Reference: FRB13460

To appear in: *Free Radical Biology and Medicine*

Received date: 31 July 2017
Accepted date: 25 September 2017

Cite this article as: Ilaria Dando, Raffaella Pacchiana, Elisa Dalla Pozza, Ivana Cataldo, Stefano Bruno, Paola Conti, Marco Cordani, Anna Grimaldi, Giovanna Butera, Michele Caraglia, Aldo Scarpa, Marta Palmieri and Massimo Donadelli, UCP2 inhibition induces ROS/Akt/mTOR axis: role of GAPDH nuclear translocation in genipin/everolimus anticancer synergism, *Free Radical Biology and Medicine*, <http://dx.doi.org/10.1016/j.freeradbiomed.2017.09.022>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

UCP2 inhibition induces ROS/Akt/mTOR axis: role of GAPDH nuclear translocation in genipin/everolimus anticancer synergism

Ilaria Dando^{1*}, Raffaella Pacchiana¹, Elisa Dalla Pozza¹, Ivana Cataldo², Stefano Bruno³, Paola Conti⁴, Marco Cordani⁵, Anna Grimaldi⁶, Giovanna Butera¹, Michele Caraglia⁶, Aldo Scarpa², Marta Palmieri¹, Massimo Donadelli^{1*}

¹Department of Neuroscience, Biomedicine and Movement, Biochemistry Section, University of Verona, Verona, Italy;

²Applied Research on Cancer Centre (ARC-Net) and Department of Diagnostics and Public Health, University of Verona, Verona, Italy;

³Food and Drug Department, University of Parma, Parma, Italy;

⁴Department of Pharmaceutical Sciences, University of Milan, Milan, Italy;

⁵Biochemistry Department; Universidad Autónoma de Madrid (UAM), Instituto de Investigaciones Biomédicas "Alberto Sols" (CSIC-UAM), IdiPAZ, Madrid, Spain;

⁶Department of Biochemistry, Biophysics and General Pathology, University of Campania "L. Vanvitelli", Naples, Italy.

massimo.donadelli@univr.it

ilaria.dando@univr.it

***Corresponding authors:** Massimo Donadelli, PhD. University of Verona, Strada Le Grazie 8, 37134 Verona, Italy. Phone: +39 045 8027281; fax: +39 045 8027170.

***Corresponding authors:** Ilaria Dando, PhD. University of Verona, Strada Le Grazie 8, 37134 Verona, Italy. Phone: +39 045 8027174; fax: +39 045 8027170.

Abstract

Several studies indicate that mitochondrial uncoupling protein 2 (UCP2) plays a pivotal role in cancer development by decreasing reactive oxygen species (ROS) produced by mitochondrial metabolism and by sustaining chemoresistance to a plethora of anticancer drugs. Here, we

Download English Version:

<https://daneshyari.com/en/article/5501616>

Download Persian Version:

<https://daneshyari.com/article/5501616>

[Daneshyari.com](https://daneshyari.com)