


Author's Accepted Manuscript

Low testosterone levels are related to oxidative stress, mitochondrial dysfunction and altered subclinical atherosclerotic markers in type 2 diabetic male patients

Susana Rovira-Llopis, Celia Bañuls, Aranzazu M de Marañon, Noelia Diaz-Morales, Ana Jover, Sandra Garzon, Milagros Rocha, Victor M. Victor, Antonio Hernandez-Mijares



www.elsevier.com

PII: S0891-5849(17)30176-4
DOI: <http://dx.doi.org/10.1016/j.freeradbiomed.2017.03.029>
Reference: FRB13271

To appear in: *Free Radical Biology and Medicine*

Received date: 3 January 2017
Revised date: 11 March 2017
Accepted date: 25 March 2017

Cite this article as: Susana Rovira-Llopis, Celia Bañuls, Aranzazu M de Marañon, Noelia Diaz-Morales, Ana Jover, Sandra Garzon, Milagros Rocha Victor M. Victor and Antonio Hernandez-Mijares, Low testosterone levels are related to oxidative stress, mitochondrial dysfunction and altered subclinical atherosclerotic markers in type 2 diabetic male patients, *Free Radical Biology and Medicine*, <http://dx.doi.org/10.1016/j.freeradbiomed.2017.03.029>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

Low testosterone levels are related to oxidative stress, mitochondrial dysfunction and altered subclinical atherosclerotic markers in type 2 diabetic male patients

Susana Rovira-Llopis^{a,1}, Celia Bañuls^{a,1}, Aranzazu M de Marañon^a, Noelia Diaz-Morales^a, Ana Jover^a, Sandra Garzon^a, Milagros Rocha^{a,b}, Victor M. Victor^{a,b,c*}, Antonio Hernandez-Mijares^{a,d*}

¹Service of Endocrinology and Nutrition, University Hospital Doctor Peset-FISABIO, Valencia, Spain

²CIBER CB06/04/0071 Research Group. CIBER Hepatic and Digestive Diseases, Department of Pharmacology, University of Valencia, Valencia, Spain

³Department of Physiology, University of Valencia, Valencia, Spain

⁴Department of Medicine, University of Valencia, Valencia, Spain

victor.victor@uv.es

hernandez_antmij@gva.es

*Corresponding authors and person to whom reprint requests should be addressed: Victor M Victor and Antonio Hernández-Mijares. Address: University Hospital Doctor Peset. Avda Gaspar Aguilar 90, 46017, Valencia, Spain. Tel.: +0034961622757, Fax: +0034961622492

Abstract

Introduction:

Low testosterone levels in men are associated with type 2 diabetes and cardiovascular risk. However, the role of testosterone in mitochondrial function and leukocyte-endothelium interactions is unknown. Our aim was to evaluate the relationship between testosterone levels,

¹ These authors have contributed equally

Download English Version:

<https://daneshyari.com/en/article/5502030>

Download Persian Version:

<https://daneshyari.com/article/5502030>

[Daneshyari.com](https://daneshyari.com)