
Accepted Manuscript

Prothrombin complex concentrate administration for bleeding associated with non-vitamin K antagonist oral anticoagulants: The SAMURAI-NVAF study

Sohei Yoshimura, Shoichiro Sato, Kenichi Todo, Yasushi Okada, Eisuke Furui, Takayuki Matsuki, Hiroshi Yamagami, Masatoshi Koga, Jun C. Takahashi, Kazuyuki Nagatsuka, Shoji Arihiro, Kazunori Toyoda, Stroke Acute Management with Urgent Riskfactor Assessment and Improvement (SAMURAI) Study Investigators



PII: S0022-510X(17)30042-4

DOI: doi: 10.1016/j.jns.2017.01.041

Reference: JNS 15103

To appear in: Journal of the Neurological Sciences

Received date: 4 September 2016 Revised date: 30 December 2016 Accepted date: 13 January 2017

Please cite this article as: Sohei Yoshimura, Shoichiro Sato, Kenichi Todo, Yasushi Okada, Eisuke Furui, Takayuki Matsuki, Hiroshi Yamagami, Masatoshi Koga, Jun C. Takahashi, Kazuyuki Nagatsuka, Shoji Arihiro, Kazunori Toyoda, Stroke Acute Management with Urgent Risk-factor Assessment and Improvement (SAMURAI) Study Investigators, Prothrombin complex concentrate administration for bleeding associated with non-vitamin K antagonist oral anticoagulants: The SAMURAI-NVAF study. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Jns(2017), doi: 10.1016/j.jns.2017.01.041

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title Page

Prothrombin Complex Concentrate Administration for Bleeding Associated with

Non-Vitamin K Antagonist Oral Anticoagulants: The SAMURAI-NVAF Study

Sohei Yoshimura, MD^a, Shoichiro Sato, MD^a, Kenichi Todo, MD^b, Yasushi Okada, MD^c, Eisuke Furui, MD^d, Takayuki Matsuki, MD^a, Hiroshi Yamagami, MD^e, Masatoshi Koga, MD^f, Jun C. Takahashi, MD^g, Kazuyuki Nagatsuka, MD^e, Shoji Arihiro, MD^a, Kazunori Toyoda, MD^a For the Stroke Acute Management with Urgent Risk-factor Assessment and Improvement (SAMURAI) Study Investigators

^a Department of Cerebrovascular Medicine, National Cerebral and Cardiovascular Center, 5-7-1, Fujishirodai, Suita-city, Osaka 565-8565, Japan

^b Department of Neurology, Kobe City Medical Center General Hospital, 2-1-1, Minatojimaminamimachi, Chuo-ku, Kobe-city, Hyogo 650-0047, Japan

^c Department of Neurology and Cerebrovascular Medicine, NHO Kyushu Medical Center, 1-8-1, Jigyohama, Chuo-ku, Fukuoka-city, Fukuoka 810-8563, Japan

^d Department of Stroke Neurology, Kohnan Hospital, 4-20-1, Nagamachiinami, Taihaku-ku, Sendai-city, Miyagi 982-8523, Japan

^e Department of Neurology, National Cerebral and Cardiovascular Center, 5-7-1, Fujishirodai, Suita-city, Osaka 565-8565, JAPAN

^f Division of Stroke Care Unit, National Cerebral and Cardiovascular Center, 5-7-1, Fujishirodai, Suita-city, Osaka 565-8565, JAPAN

^g Departments of Neurosurgery, National Cerebral and Cardiovascular Center, 5-7-1, Fujishirodai, Suita-city, Osaka 565-8565, JAPAN

Download English Version:

https://daneshyari.com/en/article/5503196

Download Persian Version:

https://daneshyari.com/article/5503196

<u>Daneshyari.com</u>