

Accepted Manuscript

Title: Access to Anti-osteoporotic Medication after Hip Fracture in Korean Elderly Patients

Authors: Yun Mi Yu, Ju-Yeun Lee, Euni Lee

PII: S0378-5122(17)30030-0

DOI: <http://dx.doi.org/doi:10.1016/j.maturitas.2017.06.021>

Reference: MAT 6841

To appear in: *Maturitas*

Received date: 16-1-2017

Revised date: 5-6-2017

Accepted date: 9-6-2017



Please cite this article as: Yu Yun Mi, Lee Ju-Yeun, Lee Euni. Access to Anti-osteoporotic Medication after Hip Fracture in Korean Elderly Patients. *Maturitas* <http://dx.doi.org/10.1016/j.maturitas.2017.06.021>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Access to Anti-osteoporosis Medication after Hip Fracture in Korean Elderly Patients

Yun Mi Yu¹, Ju-Yeun Lee², Euni Lee^{3*}

¹College of Pharmacy & Research Institute of Pharmaceutical Sciences, Seoul National University, 103 Daehak-ro, Jongno-gu, Seoul, 03080, South Korea, yupharm@snu.ac.kr

²College of Pharmacy, Institute of Pharmaceutical Science and Technology, Hanyang University, 55, Hanyangdaehak-ro, Sangrok-gu, Ansan, 15588, South Korea, jypharm@hanyang.ac.kr

³College of Pharmacy & Research Institute of Pharmaceutical Sciences, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul, 08826, South Korea, eunilee@snu.ac.kr

*Corresponding author

Euni Lee, Pharm.D. Ph.D., Associate Professor

College of Pharmacy & Research Institute of Pharmaceutical Sciences, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul, 08826, South Korea

Tel: +82-2-740-8588

Fax: 82-2-880-9122

E-mail: eunilee@snu.ac.kr

Highlights

- Access to anti-osteoporosis medication after hip fracture was found to be suboptimal in Korea.
- One in three elderly patients with hip fracture received osteoporosis drug therapy.
- Very old age, male sex, and comorbidity were risk factors for under-treatment.
- Thiazolidinedione use was a predictive factor for hip fracture treatment in women.

Download English Version:

<https://daneshyari.com/en/article/5503470>

Download Persian Version:

<https://daneshyari.com/article/5503470>

[Daneshyari.com](https://daneshyari.com)