Accepted Manuscript



Decreased linear ubiquitination of NEMO and FADD on apoptosis with caspase-mediated cleavage of HOIP

Eiji Goto, Fuminori Tokunaga

PII: S0006-291X(17)30308-X

DOI: 10.1016/j.bbrc.2017.02.040

Reference: YBBRC 37288

To appear in: *Biochemical and Biophysical Research Communications*

Received Date: 3 February 2017

Accepted Date: 7 February 2017

Please cite this article as: E. Goto, F. Tokunaga, Decreased linear ubiquitination of NEMO and FADD on apoptosis with caspase-mediated cleavage of HOIP, *Biochemical and Biophysical Research Communications* (2017), doi: 10.1016/j.bbrc.2017.02.040.

Decreased linear ubiquitination of NEMO and FADD on apoptosis with caspase-mediated cleavage of HOIP

Eiji Goto, Fuminori Tokunaga*

Department of Pathobiochemistry, Graduate School of Medicine, Osaka City University, Osaka 545-8585, Japan

*Corresponding author. Department of Pathobiochemistry, Graduate School of Medicine, Osaka City University, 1-4-3 Asahi-machi, Abeno-ku, Osaka 545-8585, Japan.

E-mail address: ftokunaga@med.osaka-cu.ac.jp (FT)