Accepted Manuscript



The membrane-tethered NAC transcription factor, AtNTL7, contributes to ER-stress resistance in *Arabidopsis*

Yong Hun Chi, Sarah Mae Boyles Melencion, Cresilda Vergara Alinapon, Min Ji Kim, Eun Seon Lee, Seol Ki Paeng, Joung Hun Park, Ganesh M. Nawkar, Young Jun Jung, Ho Byeong Chae, Chang Ho Kang, Sang Yeol Lee

PII: S0006-291X(17)30072-4

DOI: 10.1016/j.bbrc.2017.01.047

Reference: YBBRC 37109

To appear in: *Biochemical and Biophysical Research Communications*

Received Date: 9 January 2017

Accepted Date: 10 January 2017

Please cite this article as: Y.H. Chi, S.M. Boyles Melencion, C.V. Alinapon, M.J. Kim, E.S. Lee, S.K. Paeng, J.H. Park, G.M. Nawkar, Y.J. Jung, H.B. Chae, C.H. Kang, S.Y. Lee, The membrane-tethered NAC transcription factor, AtNTL7, contributes to ER-stress resistance in *Arabidopsis*, *Biochemical and Biophysical Research Communications* (2017), doi: 10.1016/j.bbrc.2017.01.047.

The membrane-tethered NAC transcription factor, AtNTL7, contributes to ER-stress resistance in *Arabidopsis*

Yong Hun Chi[a1], Sarah Mae Boyles Melencion[a1], Cresilda Vergara Alinapon[a], Min Ji Kim[a], Eun Seon Lee[a], Seol Ki Paeng[a], Joung Hun Park[a], Ganesh M. Nawkar[a], Young Jun Jung[b], Ho Byeong Chae[a], Chang Ho Kang[a], Sang Yeol Lee[a]*

[a]Division of Applied Life Science (BK21^{+} Program) and PMBBRC, Gyeongsang National University, Jinju, 52828, Korea.

[b]National Institute of Ecology, 1210 Geumgang-ro, Maseo-myeon, Seocheon-gun 33657, Republic of Korea

[1] These authors contributed equally to this work.

* **Corresponding author:** Sang Yeol Lee

Tel: +82-55-772-1351, Fax: +82-55-759-9363, E-mail: sylee@gnu.ac.kr