

# Accepted Manuscript

*Ephestia kuehniella* tolerance to *Bacillus thuringiensis* Cry1Aa is associated with reduced oligomer formation

Maissa Chakroun, Sameh Sellami, Juan Ferré, Slim Tounsi, Souad Rouis



PII: S0006-291X(16)31988-X

DOI: [10.1016/j.bbrc.2016.11.115](https://doi.org/10.1016/j.bbrc.2016.11.115)

Reference: YBBRC 36819

To appear in: *Biochemical and Biophysical Research Communications*


Received Date: 11 November 2016

Accepted Date: 21 November 2016

Please cite this article as: M. Chakroun, S. Sellami, J. Ferré, S. Tounsi, S. Rouis, *Ephestia kuehniella* tolerance to *Bacillus thuringiensis* Cry1Aa is associated with reduced oligomer formation, *Biochemical and Biophysical Research Communications* (2016), doi: 10.1016/j.bbrc.2016.11.115.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Cry1Ac is more toxic than Cry1Aa towards *E. kuehniella*.**



Download English Version:

<https://daneshyari.com/en/article/5506254>

Download Persian Version:

<https://daneshyari.com/article/5506254>

[Daneshyari.com](https://daneshyari.com)