

# Accepted Manuscript

Mouse D1Pas1, a DEAD-box RNA helicase, is required for the completion of first meiotic prophase in male germ cells

Hiroki Inoue, Narumi Ogonuki, Michiko Hirose, Yuki Hatanaka, Shogo Matoba, Shinichiro Chuma, Kimio Kobayashi, Shigeharu Wakana, Junko Noguchi, Kimiko Inoue, Kentaro Tanemura, Atsuo Ogura

PII: S0006-291X(16)31231-1

DOI: [10.1016/j.bbrc.2016.07.109](https://doi.org/10.1016/j.bbrc.2016.07.109)

Reference: YBBRC 36186

To appear in: *Biochemical and Biophysical Research Communications*

Received Date: 11 July 2016

Accepted Date: 25 July 2016

Please cite this article as: H. Inoue, N. Ogonuki, M. Hirose, Y. Hatanaka, S. Matoba, S. Chuma, K. Kobayashi, S. Wakana, J. Noguchi, K. Inoue, K. Tanemura, A. Ogura, Mouse D1Pas1, a DEAD-box RNA helicase, is required for the completion of first meiotic prophase in male germ cells, *Biochemical and Biophysical Research Communications* (2016), doi: 10.1016/j.bbrc.2016.07.109.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Mouse D1Pas1, a DEAD-box RNA helicase, is required for the  
completion of first meiotic prophase in male germ cells

Hiroki Inoue<sup>a,b,1</sup>, Narumi Ogonuki<sup>a,1</sup>, Michiko Hirose<sup>a,1</sup>, Yuki Hatanaka<sup>a,1</sup>, Shogo Matoba<sup>a</sup>,  
Shinichiro Chuma<sup>c</sup>, Kimio Kobayashi<sup>a</sup>, Shigeharu Wakana<sup>a</sup>, Junko Noguchi<sup>d</sup>, Kimiko  
Inoue<sup>a,e</sup>, Kentaro Tanemura<sup>b,\*</sup>, Atsuo Ogura<sup>a,e,f,\*</sup>

<sup>a</sup>RIKEN BioResource Center, Tsukuba, Ibaraki 305-0074, Japan

<sup>b</sup>Laboratory of Animal Reproduction and Development, Graduate School of Agricultural  
Science, Tohoku University, Miyagi 981-8555, Japan.

<sup>c</sup>Institute for Frontier Medical Sciences, Kyoto University, Kyoto 606-8507, Japan

<sup>d</sup>Institute of Agrobiological Sciences, National Agriculture and Food Research Organization  
(NARO), Tsukuba, Ibaraki 305-8602, Japan

<sup>e</sup>Graduate School of Life and Environmental Science, University of Tsukuba, Ibaraki  
305-8572, Japan

<sup>f</sup>The Center for Disease Biology and Integrative Medicine, Faculty of Medicine, University  
of Tokyo, Tokyo 113-0033, Japan

\*Corresponding author. RIKEN BioResource Center, Tsukuba, Ibaraki 305-0074, Japan and  
Laboratory of Animal Reproduction and Development, Graduate School of Agricultural  
Science, Tohoku University, Miyagi 981-8555, Japan.

*E-mail address:* ogura@rtc.riken.go.jp (A. Ogura) and kentaro@m.tohoku.ac.jp (K.  
Tanemura)

<sup>1</sup>The first four authors contributed equally to this article.

Download English Version:

<https://daneshyari.com/en/article/5506782>

Download Persian Version:

<https://daneshyari.com/article/5506782>

[Daneshyari.com](https://daneshyari.com)