



# Cationic liposomes suppress intracellular calcium ion concentration increase via inhibition of PI3 kinase pathway in mast cells



Yoshikazu Inoh<sup>a,\*</sup>, Aki Haneda<sup>a</sup>, Satoshi Tadokoro<sup>b</sup>, Satoru Yokawa<sup>a</sup>, Tadahide Furuno<sup>a</sup>

<sup>a</sup> School of Pharmacy, Aichi Gakuin University, 1-100 Kusumoto-cho, Chikusa-ku, Nagoya 464-8650, Japan

<sup>b</sup> Faculty of Pharma Sciences, Teikyo University, 2-11-1 Kaga, Itabashi-ku, Tokyo 173-8605, Japan

## ARTICLE INFO

### Keywords:

Liposome  
Allergy  
Cell signaling  
STIM1

## ABSTRACT

Cationic liposomes are commonly used as vectors to effectively introduce foreign genes (antisense DNA, plasmid DNA, siRNA, etc.) into target cells. Cationic liposomes are also known to affect cellular immunocompetences such as the mast cell function in allergic reactions. In particular, we previously showed that the cationic liposomes bound to the mast cell surface suppress the degranulation induced by cross-linking of high affinity IgE receptors in a time- and dose-dependent manner. This suppression is mediated by impairment of the sustained level of intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) via inhibition of store-operated  $\text{Ca}^{2+}$  entry (SOCE). Here we study the mechanism underlying an impaired  $[\text{Ca}^{2+}]_i$  increase by cationic liposomes in mast cells. We show that cationic liposomes inhibit the phosphorylation of Akt and PI3 kinases but not Syk and LAT. As a consequence, SOCE is suppressed but  $\text{Ca}^{2+}$  release from endoplasmic reticulum (ER) is not. Cationic liposomes inhibit the formation of STIM1 puncta, which is essential to SOCE by interacting with Orai1 following the  $\text{Ca}^{2+}$  concentration decrease in the ER. These data suggest that cationic liposomes suppress SOCE by inhibiting the phosphorylation of PI3 and Akt kinases in mast cells.

## 1. Introduction

Several studies have shown that cationic liposomes exert immunomodulatory effects associated with low immunogenicity and toxicity and offer advantages such as easy preparation and targeting [1–4]. Cationic liposomes not only transport DNA to immune cells but also enhance the function of antigen-presenting cells such as dendritic cells and macrophages [5–7]. Previously, we investigated the effect of a particular cationic liposome on the mast cell function during allergic reactions and found that the cationic liposomes suppressed degranulation induced by cross-linking of high affinity IgE receptors (FcεRI) in a time- and dose-dependent manner [8]. The suppression of degranulation is mediated by the impairment of a sustained level of intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) derived from the inhibition of store-operated  $\text{Ca}^{2+}$  entry (SOCE). Furthermore, cationic liposomes suppress vascular permeability elevation induced by mast cell activation in mice.

Mast cells play important roles in innate and adaptive immune responses [9]. Cross-linking of FcεRI on mast cells by multivalent antigens induces two pathways activated by src-family protein tyrosine kinases, Lyn and Fyn [10]. Lyn phosphorylates the tyrosine residues in the immunoreceptor tyrosine-based activation motif (ITAM) of FcεRI β- and γ-chains. Phosphorylated ITAMs of the γ-chain create binding sites for the

Src homology 2 (SH2) domains of the tyrosine kinase Syk, which is then activated by phosphorylation. The activated Syk subsequently phosphorylates linker for activation of T cells (LAT), leading to the organization of a complex with phospholipase C (PLC) γ [11]. Activation of PLCγ in this complex causes the hydrolysis of phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>) into inositol-1,4,5-triphosphate (IP<sub>3</sub>) and diacylglycerol (DAG), increasing the  $[\text{Ca}^{2+}]_i$  through  $\text{Ca}^{2+}$  release from endoplasmic reticulum (ER) [12]. This pathway is known as the Lyn–Syk–LAT pathway. The LAT-activating pathway is the main contributor to  $\text{Ca}^{2+}$  depletion in the ER because LAT organizes the complexes comprising PLCγ, SLP76, and other adaptor proteins.

In addition to Syk, the second src-family protein tyrosine kinase Fyn also phosphorylates Grb-associated binder 2 (Gab2), an adapter protein essential for phosphatidylinositol-3 kinase–Akt activation (PI3K–Akt pathway) [13–15]. The activation of this pathway leads to extracellular  $\text{Ca}^{2+}$  influx, an indispensable step for mast cell degranulation. To regulate SOCE following a  $\text{Ca}^{2+}$  decrease in ER, the ER protein stromal interaction molecule 1 (STIM1) forms oligomers, translocates to the plasma membrane region (ER–PM junctions), and interacts with Orai1 [16,17]. Indeed, ER–PM junctions are often visualized as the formation of STIM1 puncta [18,19].

The aim of this study was to examine the mechanism of impaired

\* Corresponding author.

E-mail address: [inoh@dpc.agu.ac.jp](mailto:inoh@dpc.agu.ac.jp) (Y. Inoh).

<http://dx.doi.org/10.1016/j.bbamem.2017.09.025>

Received 14 June 2017; Received in revised form 7 September 2017; Accepted 27 September 2017

Available online 29 September 2017

0005-2736/© 2017 Elsevier B.V. All rights reserved.

$[Ca^{2+}]_i$  increase by the treatment with cationic liposomes in RBL-2H3 cells. In this paper, we report that cationic liposomes suppress SOCE through the inhibition of PI3K and Akt activation. Our results might contribute to the clinical development of cationic liposomes as a useful therapeutic tool for allergic diseases mediated by mast cell activation.

## 2. Materials and methods

### 2.1. Materials

The synthesis of the cationic cholesterol derivative containing a hydrophilic amino head group, cholesteryl-3 $\beta$ -carboxyamidoethylene-N-hydroxyethylamine (OH-Chol) was described in a previous study [20] and 1,2-dioleoyl-*sn*-glycero-3-phosphatidylethanolamine (DOPE) was purchased from Sigma-Aldrich (St. Louis, MO, USA). Mouse anti-dinitrophenyl (DNP)-IgE was provided by Prof. T. Kishimoto (Osaka University). DNP-conjugated bovine serum albumin (DNP-BSA) was prepared using previously described methods [21]. In the present experiments, six DNP groups, on average, were conjugated with a single BSA molecule. Thapsigargin was purchased from Wako Pure Chemical Industries (Osaka, Japan). PhosphoBLOCKER™, rabbit anti-phospho-Syk(Y323) antibody, rabbit anti-phospho-LAT(Y191) antibody, rabbit anti-phospho-PI3K p85(Y458)/p55(Y199) antibody, rabbit anti-phospho-Akt(T308) antibody, horse radish peroxidase (HRP)-labeled goat anti-rabbit IgG, HRP-labeled goat anti-mouse IgG, and rabbit anti-STIM1 antibodies were purchased from Cell Signaling Technology (Danvers, MA, USA). Rabbit anti-Orai1 antibody was purchased from Proteintech (Rosemont, IL, USA). Mouse anti- $\beta$ -actin antibody was purchased from Sigma-Aldrich. Alexa Fluor 555-labeled goat anti-rabbit IgG antibody was obtained from Molecular Probes (Cambridge Cambridgeshire, UK).

### 2.2. Preparation of liposomes

The cationic liposomes were prepared by ultrasonication methods according to our previously described procedures [22–24]. In brief, DOPE (20 nmol) and OH-Chol (30 nmol) were mixed and dried with  $N_2$  gas under reduced pressure to remove the chloroform solvent. The lipid film was hydrated with 400  $\mu$ l of phosphate buffered saline (PBS) for 1 h. Samples were sonicated in a bath-type sonicator (Branson model B 1200) to generate small unilamellar vesicles. We measured the average size of cationic liposome by dynamic light scattering using zetasizer Nano ZS90 (Malvern Instruments, Worcestershire, UK) with a scattering angle of 90°. The size of cationic liposome was  $453.2 \pm 83.2$  nm. We confirmed that the size peak of liposome and lipoplexes was so sharp and polydispersity index was low.

### 2.3. Cell culture

The rat mast cell line RBL-2H3, which is the most widely used, as a model for mast cells, was cultured in minimum essential medium (Nissui, Tokyo, Japan) supplemented with 10% fetal calf serum (Roche, Mannheim, Germany).

### 2.4. Western blot analysis

Cultured RBL-2H3 cells were seeded in 60 mm plates at a density of  $5 \times 10^5$  cells/well 24 h before introduction of the cationic liposomes. Cells were incubated with 23.3  $\mu$ g/ml of cationic liposomes for 3 h and washed in HEPES buffer [10 mM HEPES (pH 7.2), 140 mM NaCl, 5 mM KCl, 0.6 mM  $MgCl_2$ , 1 mM  $CaCl_2$ , 0.1% glucose, 0.1% BSA, and 0.01% sulfapyrazone]. After sensitizing the RBL-2H3 cells with anti-DNP IgE for 30 min at 37 °C, the cells were washed twice with HEPES buffer. Thereafter, cells were lysed with cold lysis buffer [20 mM HEPES (pH 7.9), 0.1% NP-40, 50 mM NaCl, 1 mM EDTA, 1 mM dithiothreitol, 10 mM  $Na_3VO_4$ , 10  $\mu$ g/ml phenylmethylsulfonyl fluoride, 10  $\mu$ g/ml

leupeptin, and 10% glycerol]. The lysates were kept on ice for 30 min, followed by centrifugation at 15,000 rpm for 20 min at 0 °C. The resulting supernatants were solubilized by treatment with Laemmli buffer at 100 °C for 3 min. Subsequently, proteins were separated using 10% SDS-polyacrylamide gel electrophoresis. The separated proteins were transferred to a PVDF membrane with an electroblotter. After blocking with 5% PhosphoBLOCKER™, the membranes were probed with rabbit anti-phospho-Syk (Y323) (1:1000), rabbit anti-phospho-LAT (Y191) (1:1000), rabbit anti-phospho-PI3K p85 (Y458)/p55(Y199) (1:1000), rabbit anti-phospho-Akt (T308) (1:1000), rabbit anti-STIM1 (1:1000), mouse anti-Orai1 (1:1000), or mouse anti- $\beta$ -actin (1:40,000) antibody at 4 °C overnight. Subsequently, treatment with HRP-labeled goat anti-rabbit IgG (1:2000) or HRP-labeled goat anti-mouse IgG (1:2000) was performed. Immunoreactivity was detected using enhanced chemiluminescence (ECL; GE Healthcare, Buckinghamshire, UK) with a LAS-3000mini (Fujifilm, Tokyo, Japan) and analyzed using Image Gauge (Fujifilm).

### 2.5. Confocal laser scanning microscopy (CLSM)

RBL-2H3 cells were incubated with 23.3  $\mu$ g/ml cationic liposomes for 3 h and washed in HEPES buffer. After sensitizing the RBL-2H3 cells with anti-DNP IgE at 37 °C, cells were washed twice with HEPES buffer and then immediately fixed with ice-cold paraformaldehyde in PBS overnight. The next day, cells were incubated with rabbit anti-STIM1 (1:800) at 4 °C overnight. Subsequently, treatment with Alexa Fluor 555-labeled goat anti-rabbit IgG antibody (1:400) was performed. The distribution of STIM1 was analyzed in an observation chamber (Elecon, Chiba, Japan) by CLSM (LSM 510-META; Zeiss, Oberkochen, Germany). The Alexa Fluor 555-labeled goat anti-rabbit IgG antibody was excited by a He-Ne laser (543 nm), and fluorescence was detected through a band-pass filter at 560–580 nm.

### 2.6. Measurement of $[Ca^{2+}]_i$

RBL-2H3 cells ( $5 \times 10^5$  cells) were loaded with 1  $\mu$ M Fura 2-AM (Molecular Probes, Eugene, OR, USA.), sensitized with anti-DNP IgE for 30 min at 37 °C, and then washed twice with HEPES buffer or  $Ca^{2+}$ -free HEPES buffer. Fluorescence intensities at 500 nm using excitation wavelengths of 340 and 360 nm (F340 and F360, respectively) were measured at 37 °C, and the F340/F360 ratio was calculated using a spectrofluorometer connected to a personal computer (RF-5300PC; Shimadzu, Kyoto, Japan). Ratios were converted to  $Ca^{2+}$  concentrations by a previously described procedure [25].

## 3. Results

### 3.1. Effects of cationic liposomes on $Ca^{2+}$ release from ER and SOCE

Initially, to check the effects of cationic liposomes on  $Ca^{2+}$  release from ER, we measured the  $[Ca^{2+}]_i$  in RBL-2H3 cells in the absence of extracellular  $Ca^{2+}$  following antigen stimulation. We found that there was no significant difference in the  $[Ca^{2+}]_i$  increase in RBL-2H3 cells with and without treatment with cationic liposomes (22.3  $\mu$ g/ml) for 3 h (Fig. 1), indicating that cationic liposomes do not affect the amount of  $Ca^{2+}$  release from ER. On the contrary, the pretreatment with cationic liposomes significantly suppressed the  $[Ca^{2+}]_i$  increase in the presence of extracellular  $Ca^{2+}$  (3 mM) (Fig. 1). This result indicates that cationic liposomes inhibit SOCE but not  $Ca^{2+}$  release from ER in RBL-2H3 cells stimulated by antigen.

### 3.2. Effects of cationic liposomes on phosphorylation of signaling proteins

In mast cells, when Fc $\epsilon$ RI are crosslinked by multivalent antigens, two pathways are activated via src-kinase Lyn and Fyn. Although both Lyn and Fyn phosphorylate Syk, Lyn and Fyn subsequently activate LAT

Download English Version:

<https://daneshyari.com/en/article/5507314>

Download Persian Version:

<https://daneshyari.com/article/5507314>

[Daneshyari.com](https://daneshyari.com)