

Cyp7a1 is continuously increased with disrupted Fxr-mediated feedback inhibition in hypercholesterolemic TALLYHO/Jng mice

Eun-Ah Lee, Dong-In Lee, Hee-Yoen Kim, Sung-Hoon Ahn, Hye-Rim Seong, Won Hoon Jung, Ki Young Kim, Sungsub Kim, Sang Dal Rhee

PII: S1388-1981(17)30174-9
DOI: doi:10.1016/j.bbalip.2017.08.007
Reference: BBAMCB 58196

To appear in: *BBA - Molecular and Cell Biology of Lipids*

Received date: 21 February 2017
Revised date: 18 August 2017
Accepted date: 22 August 2017

Please cite this article as: Eun-Ah Lee, Dong-In Lee, Hee-Yoen Kim, Sung-Hoon Ahn, Hye-Rim Seong, Won Hoon Jung, Ki Young Kim, Sungsub Kim, Sang Dal Rhee, Cyp7a1 is continuously increased with disrupted Fxr-mediated feedback inhibition in hypercholesterolemic TALLYHO/Jng mice, *BBA - Molecular and Cell Biology of Lipids* (2017), doi:10.1016/j.bbalip.2017.08.007

Cyp7a1 is continuously increased with disrupted Fxr-mediated feedback inhibition in hypercholesterolemic TALLYHO/Jng mice

Eun-Ah Lee[1,2], Dong-In Lee[1,2], Hee-Yoen Kim[1], Sung-Hoon Ahn[3], Hye-Rim Seong[1], Won Hoon Jung[1], Ki Young Kim[1,2], Sungsub Kim[2], Sang Dal Rhee[1,2,*]

1. Bio & Drug Discovery Division, Korea Research Institute of Chemical Technology, 141, Gajeong-ro, Yuseong-gu, Daejeon, 34114, Korea

2. Graduate School of New Drug Discovery and Development, Chungnam National University, 99, Daehak-ro, Yuseong-gu, Daejeon, 34134, Korea

3. College of Pharmacy, Kangwon National University, Chuncheon, 24341, Korea

*Corresponding author

Bio & Drug Discovery Division, Korea Research Institute of Chemical Technology, 141, Gajeong-ro, Yuseong-gu, Daejeon, 34114, Korea

Tel.: +82-042-860-7421, Fax: +82-042-862-0770

E-mail: sdrhee@krict.re.kr