

Accepted Manuscript

An aromatic amino acid within intracellular loop 2 of the prostaglandin EP2 receptor is a prerequisite for selective association and activation of G α s

Akiko Yano, Yuko Takahashi, Hiromi Moriguchi, Tomoaki Inazumi, Tomoaki Koga, Akira Otaka, Yukihiro Sugimoto

PII: S1388-1981(17)30054-9
DOI: doi:[10.1016/j.bbalip.2017.03.006](https://doi.org/10.1016/j.bbalip.2017.03.006)
Reference: BBAMCB 58129

To appear in: *BBA - Molecular and Cell Biology of Lipids*

Received date: 13 September 2016
Revised date: 17 March 2017
Accepted date: 18 March 2017



Please cite this article as: Akiko Yano, Yuko Takahashi, Hiromi Moriguchi, Tomoaki Inazumi, Tomoaki Koga, Akira Otaka, Yukihiro Sugimoto, An aromatic amino acid within intracellular loop 2 of the prostaglandin EP2 receptor is a prerequisite for selective association and activation of G α s, *BBA - Molecular and Cell Biology of Lipids* (2017), doi:[10.1016/j.bbalip.2017.03.006](https://doi.org/10.1016/j.bbalip.2017.03.006)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

MS#: BBALIP-16-269 3rd Revision

An aromatic amino acid within intracellular loop 2 of the prostaglandin EP2 receptor is a prerequisite for selective association and activation of G α s

Akiko Yano^a, Yuko Takahashi^a, Hiromi Moriguchi^a, Tomoaki Inazumi^{b,c}, Tomoaki Koga^d, Akira Otaka^e, and Yukihiro Sugimoto^{a,b,c,*}

^aDepartment of Physiological Chemistry, Graduate School of Pharmaceutical Sciences, Kyoto University, Kyoto, Japan

^bDepartment of Pharmaceutical Biochemistry, Graduate School of Pharmaceutical Sciences, Kumamoto University, Kumamoto, Japan

^cAMED-CREST, Tokyo 100-0004, Japan

^dPriority Organization for Innovation and Excellence, Kumamoto University, Kumamoto, Japan

^eDepartment of Bioorganic Synthetic Chemistry, Graduate School of Pharmaceutical Sciences, Tokushima University, Tokushima, Japan

*Corresponding author at: Department of Pharmaceutical Biochemistry, Graduate School of Pharmaceutical Sciences, Kumamoto University, Oe-honmachi, Chuo-ku, Kumamoto 862-0973, Japan. E-mail: ysugi@kumamoto-u.ac.jp (Yukihiro Sugimoto)

Keywords: eicosanoid, G protein-coupled receptor, heterotrimeric G protein, prostaglandin, prostanoid receptor, receptor structure-function

Download English Version:

<https://daneshyari.com/en/article/5508432>

Download Persian Version:

<https://daneshyari.com/article/5508432>

[Daneshyari.com](https://daneshyari.com)