

Cold-sensing TRPM8 channel participates in circadian control of the brown adipose tissue

Maria Nathália Moraes, Leonardo Vinicius Monteiro de Assis, Felipe dos Santos Henriques, Miguel Luiz Batista Junior, Ali D. Güler, Ana Maria de Lauro Castrucci

PII: S0167-4889(17)30254-9
DOI: doi:10.1016/j.bbamcr.2017.09.011
Reference: BBAMCR 18179

To appear in:

Received date: 5 June 2017
Revised date: 18 September 2017
Accepted date: 19 September 2017

Please cite this article as: Maria Nathália Moraes, Leonardo Vinicius Monteiro de Assis, Felipe dos Santos Henriques, Miguel Luiz Batista Junior, Ali D. Güler, Ana Maria de Lauro Castrucci , Cold-sensing TRPM8 channel participates in circadian control of the brown adipose tissue. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Bbamcr(2017), doi:10.1016/j.bbamcr.2017.09.011

Cold-sensing TRPM8 channel participates in circadian control of the brown adipose tissue

Maria Nathália Moraes[1], Leonardo Vinicius Monteiro de Assis[1], Felipe dos Santos Henriques[2,3], Miguel Luiz Batista Junior[3], Ali D. Güler[4], Ana Maria de Lauro Castrucci[1,4]

[1] Institute of Biosciences, Department of Physiology, University of São Paulo, São Paulo, Brazil;

[2] Program in Molecular Medicine, University of Massachusetts Medical School, Worcester, Massachusetts, USA;

[3] Laboratory of Adipose Tissue Biology, Integrated Group of Biotechnology, University of Mogi das Cruzes, Mogi das Cruzes, Brazil;

[4] Department of Biology, University of Virginia, Charlottesville, VA, USA.

Corresponding author: Ana Maria de Lauro Castrucci, Departamento de Fisiologia, Instituto de Biociências, Universidade de São Paulo, R. do Matão, trav. 14, no. 101, São Paulo, 05508-900, Brazil; email: amdlcast@ib.usp.br

Running title: TRPM8 channel and peripheral tissue clock

Key words: *Mus musculus*, suprachiasmatic nucleus, eye, brown adipose tissue, TRPM8 channel, clock gene machinery