

HPI/AMF inhibition halts the development of the aggressive phenotype of breast cancer stem cells

Juan Carlos Gallardo-Pérez, Alhelí Adán-Ladrón de Guevara, Alvaro Marín-Hernández, Rafael Moreno-Sánchez, Sara Rodríguez-Enríquez

PII: S0167-4889(17)30167-2
DOI: doi:10.1016/j.bbamcr.2017.06.015
Reference: BBAMCR 18128

To appear in: *BBA - Molecular Cell Research*

Received date: 2 January 2017
Revised date: 13 June 2017
Accepted date: 16 June 2017

Please cite this article as: Juan Carlos Gallardo-Pérez, Alhelí Adán-Ladrón de Guevara, Alvaro Marín-Hernández, Rafael Moreno-Sánchez, Sara Rodríguez-Enríquez, HPI/AMF inhibition halts the development of the aggressive phenotype of breast cancer stem cells, *BBA - Molecular Cell Research* (2017), doi:10.1016/j.bbamcr.2017.06.015

HPI/AMF inhibition halts the development of the aggressive phenotype of breast cancer stem cells

Juan Carlos Gallardo-Pérez*, Alhelí Adán-Ladrón de Guevara, Alvaro Marín-Hernández, Rafael Moreno-Sánchez and Sara Rodríguez-Enríquez*

Departamento de Bioquímica, Instituto Nacional de Cardiología, Tlalpan, México, DF 14080, México.

Authors for correspondence:

*Sara Rodríguez-Enríquez, Ph. D. and
Juan Carlos Gallardo-Pérez, Ph. D.
Instituto Nacional de Cardiología
Departamento de Bioquímica
Juan Badiano No. 1, Col. Sección XVI
Tlalpan, México 14080
MEXICO
saren960104@hotmail.com

Running Title: HPI/AMF inhibition blocks cancer stem cells phenotype

Keywords: Breast cancer; stem cells; hexose-phosphate isomerase; metastatic phenotype

Abbreviations: BCSC, breast cancer stem cell-like cells; CSC, cancer stem cells; HPI/AMF, hexose phosphate isomerase/autocrine motility factor; E4P, erythrose 4 phosphate; EMT, epithelial mesenchymal transition.