Accepted Manuscript

Prevalence of anti-NT5C1A antibodies in Japanese patients with autoimmune rheumatic diseases in comparison with other patient cohorts

Yoshinao Muro, Hirotaka Nakanishi, Masahisa Katsuno, Michihiro Kono, Masashi Akiyama


PII: S0009-8981(17)30252-8

DOI: doi: 10.1016/j.cca.2017.07.002

Reference: CCA 14790

To appear in: Clinica Chimica Acta

Received date: 8 June 2017 Revised date: 30 June 2017 Accepted date: 3 July 2017



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Full-length research article

Prevalence of anti-NT5C1A antibodies in Japanese patients

with autoimmune rheumatic diseases

in comparison with other patient cohorts

Yoshinao Muro^a, Hirotaka Nakanishi^b, Masahisa Katsuno^b,

Michihiro Kono^a, Masashi Akiyama^a

^aDepartment of Dermatology, Nagoya University Graduate School of Medicine, Nagoya,

Japan

^bDepartment of Neurology, Nagoya University Graduate School of Medicine, Nagoya, Japan

Correspondence to: Yoshinao Muro, MD, PhD

Division of Connective Tissue Disease and Autoimmunity, Department of Dermatology,

Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku, Nagoya

466-8550, JAPAN

TEL: 81-52-744-2314

FAX: 81-52-744-2318

E-mail: ymuro@med.nagoya-u.ac.jp

Download English Version:

https://daneshyari.com/en/article/5509581

Download Persian Version:

https://daneshyari.com/article/5509581

Daneshyari.com