

Accepted Manuscript

Sardine procalcitonin amino-terminal cleavage peptide has a different action from calcitonin and promotes osteoblastic activity in the scales of goldfish

Yoichi Kase, Takahiro Ikari, Toshio Sekiguchi, Masayuki Sato, Shouzo Ogiso, Tsuyoshi Kawada, Shin Matsubara, Honoo Satake, Yuichi Sasayama, Masato Endo, Kei-ichiro Kitamura, Atsuhiko Hattori, Takushi X. Watanabe, Yusuke Maruyama, Yoshinari Watanabe, Hisayuki Funahashi, Akira Kambe-gawa, Nobuo Suzuki

PII: S1095-6433(17)30135-6
DOI: doi:[10.1016/j.cbpa.2017.06.007](https://doi.org/10.1016/j.cbpa.2017.06.007)
Reference: CBA 10239

To appear in: *Comparative Biochemistry and Physiology, Part A*

Received date: 18 March 2017
Revised date: 31 May 2017
Accepted date: 7 June 2017

Please cite this article as: Kase, Yoichi, Ikari, Takahiro, Sekiguchi, Toshio, Sato, Masayuki, Ogiso, Shouzo, Kawada, Tsuyoshi, Matsubara, Shin, Satake, Honoo, Sasayama, Yuichi, Endo, Masato, Kitamura, Kei-ichiro, Hattori, Atsuhiko, Watanabe, Takushi X., Maruyama, Yusuke, Watanabe, Yoshinari, Funahashi, Hisayuki, Kambe-gawa, Akira, Suzuki, Nobuo, Sardine procalcitonin amino-terminal cleavage peptide has a different action from calcitonin and promotes osteoblastic activity in the scales of goldfish, *Comparative Biochemistry and Physiology, Part A* (2017), doi:[10.1016/j.cbpa.2017.06.007](https://doi.org/10.1016/j.cbpa.2017.06.007)



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Sardine prolactin amino-terminal cleavage peptide has a different action from calcitonin and promotes osteoblastic activity in the scales of goldfish

Yoichi Kase¹, Takahiro Ikari¹, Toshio Sekiguchi¹, Masayuki Sato¹, Shouzo Ogiso¹, Tsuyoshi Kawada², Shin Matsubara², Honoo Satake², Yuichi Sasayama¹, Masato Endo³, Kei-ichiro Kitamura⁴, Atsuhiko Hattori⁵, Takushi X. Watanabe⁶, Yusuke Maruyama⁵, Yoshinari Watanabe⁷, Hisayuki Funahashi⁸, Akira Kambegawa⁹ and Nobuo Suzuki^{1*}

¹Noto Marine Laboratory, Institute of Nature and Environmental Technology, Kanazawa University, Noto-cho, Ishikawa 927-0553, Japan

²Bioorganic Research Institute, Suntory Foundation for Life Sciences, 8-1-1, Seikadai, Seika-cho, Soraku-gun, Kyoto 619-0284, Japan

³Graduate School of Marine Science and Technology, Tokyo University of Marine Science and Technology, Minato-ku, Tokyo 108-8477, Japan

⁴Department of Clinical Laboratory Science, Graduate School of Medical Science, Kanazawa University, Kanazawa, Ishikawa 920-0942, Japan

⁵Department of Biology, College of Liberal Arts and Sciences, Tokyo Medical and Dental University, Ichikawa, Chiba 272-0827, Japan

⁶Peptide Institute Inc., Osaka 567-0085, Japan

⁷Organization of Frontier Science and Innovation, Kanazawa University, Kakuma, Ishikawa 920-1192, Japan

⁸Department of Anatomy, Showa University School of Medicine, Shinagawa-ku, Tokyo 142-8555, Japan

⁹Kambegawa Laboratory, Tokyo 201-0013, Japan

*Correspondence to: Dr. Nobuo. Suzuki,
Noto Marine Laboratory, Institute of Nature and Environmental Technology,
Kanazawa University, Ogi, Noto-cho, Ishikawa 927-0553, Japan.
Tel.: 81-768-74-1151; Fax 81-768-74-1644;
e-mail: nobuos@staff.kanazawa-u.ac.jp

Download English Version:

<https://daneshyari.com/en/article/5510276>

Download Persian Version:

<https://daneshyari.com/article/5510276>

[Daneshyari.com](https://daneshyari.com)