

Accepted Manuscript

Title: NEUROACTIVE STEROID LEVELS AND PSYCHIATRIC AND ANDROLOGICAL FEATURES IN POST-FINASTERIDE PATIENTS.

Authors: Roberto Cosimo Melcangi, Daniele Santi, Roberto Spezzano, Maria Grimoldi, Tommaso Tabacchi, Maria Letizia Fusco, Silvia Diviccaro, Silvia Giatti, Giuseppe Carrà, Donatella Caruso, Manuela Simoni, Guido Cavaletti



PII: S0960-0760(17)30102-4
DOI: <http://dx.doi.org/doi:10.1016/j.jsbmb.2017.04.003>
Reference: SBMB 4926

To appear in: *Journal of Steroid Biochemistry & Molecular Biology*

Received date: 30-12-2016
Revised date: 22-2-2017
Accepted date: 6-4-2017

Please cite this article as: Roberto Cosimo Melcangi, Daniele Santi, Roberto Spezzano, Maria Grimoldi, Tommaso Tabacchi, Maria Letizia Fusco, Silvia Diviccaro, Silvia Giatti, Giuseppe Carrà, Donatella Caruso, Manuela Simoni, Guido Cavaletti, NEUROACTIVE STEROID LEVELS AND PSYCHIATRIC AND ANDROLOGICAL FEATURES IN POST-FINASTERIDE PATIENTS., *Journal of Steroid Biochemistry and Molecular Biology* <http://dx.doi.org/10.1016/j.jsbmb.2017.04.003>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

HIGHLIGHTS

- **In patients treated with finasteride for male pattern hair loss, persistent side effects may occur**
- **Erectile dysfunction and abnormal somatosensory evoked potentials of the pudendal nerve were reported**
- **Major depressive disorder and altered levels of neuroactive steroids were observed.**

Download English Version:

<https://daneshyari.com/en/article/5513300>

Download Persian Version:

<https://daneshyari.com/article/5513300>

[Daneshyari.com](https://daneshyari.com)