



ELSEVIER

Contents lists available at ScienceDirect

Journal of Bone Oncology

journal homepage: www.elsevier.com/locate/jbo

Research paper

Surgical management of proximal fibular tumors: A report of 12 cases



Hiroyuki Inatani ^{a,b}, Norio Yamamoto ^a, Katsuhiko Hayashi ^{a,b,*}, Hiroaki Kimura ^{a,b}, Akihiko Takeuchi ^a, Shinji Miwa ^a, Takashi Higuchi ^a, Kensaku Abe ^a, Yuta Taniguchi ^a, Satoshi Yamada ^b, Hideki Okamoto ^b, Takanobu Otsuka ^b, Hiroyuki Tsuchiya ^a

^a Department of Orthopaedic Surgery, Kanazawa University Graduate School of Medical Science, 13-1, Takara-machi, Kanazawa-shi, Ishikawa-ken 920-8640, Japan

^b Department of Orthopaedic Surgery, Nagoya City University Graduate School of Medical Science, 1, Kawasumi, Mizuho-cho, Mizuho-ku, Nagoya-shi, Aichi-ken 467-8601, Japan

ARTICLE INFO

Article history:

Received 30 May 2016

Accepted 4 June 2016

Available online 8 June 2016

Keywords:

Proximal fibular tumor

Reconstruction

Fibular head resection

ABSTRACT

Background/aim: Aggressive benign or malignant tumors in the proximal fibula may require en bloc resection of the fibular head, including the peroneal nerve and lateral collateral ligament. Here, we report the treatment outcomes of 12 patients with aggressive benign or malignant proximal fibula tumors.

Patients and methods: Four patients with osteosarcoma and 1 patient with Ewing's sarcoma were treated with intentional marginal resections after effective chemotherapy, and 4 patients underwent fibular head resections without ligamentous reconstruction. Clinical outcomes were investigated.

Results: The mean Musculoskeletal Tumor Society scores were 96% and 65% in patients without peroneal nerve resection and those with nerve resection, respectively. No patients complained of knee instability.

Conclusion: Functional outcomes after resection of the fibular head were primarily influenced by peroneal nerve preservation. If patients are good responders to preoperative chemotherapy, malignant tumors may be treated with marginal excision, resulting in peroneal nerve preservation and good function.

© 2016 The Authors. Published by Elsevier GmbH. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Osteosarcomas (OS), giant cell tumors (GCT), and chondrosarcomas (CS) rarely arise in the proximal fibula, as tumors of the fibula account for only 2.5% of all primary bone tumors [1,2]. Approximately one-fourth of all primary bone tumors in the fibula are malignant [2], and approximately one-fourth of all benign bone tumors in the fibula are GCT. Aggressive benign tumors such as GCT and malignant tumors in the proximal fibula often require en bloc resection of the fibular head, including the lateral collateral ligament (LCL) and the attachment site of the biceps femoris muscle tendon. Fibular head resection may cause postoperative knee instability because the LCL is the main resistor of varus loading. However, unlike traumatic disruption of the lateral structures of the knee, the need for reconstruction of the LCL after tumor excision is controversial. The majority of traumatic posterolateral knee injuries occur in combination with other ligamentous injuries [3], the most common of which affect the

anterior cruciate ligament or posterior cruciate ligament [4]. During proximal fibular tumor excision, these ligaments are often preserved.

Peroneal nerve palsy and local recurrence are serious postoperative complications associated with resection of these tumors. Based on the literature, the incidence rate of postoperative peroneal nerve palsy ranges from 3% to 57% [5], while local recurrence rates vary by tumor histology and resection type.

In this study, we investigated the clinical outcomes of 12 patients with aggressive benign or malignant proximal fibula tumors, especially in terms of the necessity for LCL reconstruction and the impact of peroneal nerve resection on postoperative function.

2. Patients and methods

We retrospectively reviewed our institution's medical records to identify all patients with tumors of the proximal fibula surgically treated between 1992 and 2011. Nonoperative cases were excluded. The proximal epiphysis was involved in all patients. The clinical outcomes of all patients were investigated. All patients approved the use of their data for this study, and this study was approved by the ethics committee at our institution.

* Corresponding author at: Department of Orthopaedic Surgery, Kanazawa University Graduate School of Medical Science, 13-1, Takara-machi, Kanazawa-shi, Ishikawa-ken 920-8640, Japan.

E-mail address: hayashikatsu830@aol.com (K. Hayashi).

We surgically treated 12 patients (7 men, 5 women). The mean age at presentation was 31 years (range, 11–71 years). Preoperative metastatic workups were performed by chest X-ray in all patients, and especially by chest CT in all patients of malignancy. Patients were followed up for a minimum of 2.7 years (median, 5.2 years; range, 2.7–13.4 years) at intervals of 2 weeks, 3 months, 6 months, 1 year, 2 years, 3 years, 4 years, and 5 years postoperatively. Follow-up intervals and durations were determined individually for each patient and according to tumor pathology.

Presenting symptoms were available for 10 patients, and these included pain (9/10), a palpable mass (1/10), and peroneal nerve symptoms (3/10).

Histological diagnoses were OS (4/12), malignant fibrous histiocytoma (MFH; 2/12), Ewing's sarcoma (EWS; 1/12), GCT (4/12), and low-grade CS (1/12).

We analyzed surgical methods, oncologic outcomes, functional results, and complication rates among patients. Surgical methods were classified into 3 types: intralesional excision, marginal excision, and wide excision (Fig. 1). Functional results were assessed using Musculoskeletal Tumor Society scores, in which numerical values (scores ranging from 0 to 5, with 0 indicating poor function and 30 indicating good function) are assigned to each of 6 categories: pain, function, emotional acceptance, support, walking, and gait. Data were expressed as a percentage of total scores between 0 and 30.

3. Results

Intralesional excision was performed in 4 GCT patients by curettage and adjuvant therapy with burring, cryotherapy, and alcohol. Intentional marginal excision was performed in 4 OS patients and 1 EWS patient following effective caffeine-potentiated chemotherapy in which the histological responses of resected specimens were classified as Grade IV (Rosen and Huvos) [6]. One patient with low-grade CS without extraskelatal lesions and 2 patients with recurrent GCT were also treated with marginal excision of the fibular head. Wide excision, including peroneal nerve resection, was performed for 2 MFH patients (Table 1).

Ten patients, including 2 with recurrent GCT, were treated by fibular head resection. Four patients, including 2 MFH patients, 1 EWS patient, and 1 recurrent GCT patient, were treated by fibular head resection without ligamentous reconstruction. In the other 6 patients who underwent fibular head resection, the LCL and biceps tendon were reattached to the lateral tibia with suture anchors, a spike washer, or just sutures.

At a median follow-up of 5.2 years, 8 patients were disease-free (median follow-up, 7.7 years), 2 patients had no evidence of disease (median follow-up, 4.4 years), 1 patient was alive with disease, and 1 OS patient died of disease. The OS patient never had local recurrence though metastatic tumors were detected 2 years after first operation.

The mean Musculoskeletal Tumor Society score in patients without peroneal nerve resection was 96% (range, 70–100%), while that in patients with peroneal nerve resection was 65% (range, 60–70%). Permanent peroneal nerve palsy, despite nerve preservation, occurred in 1 of 10 patients. Deep venous thrombosis (1/12) also occurred in the same patient during a period of bed rest following neurolysis; therefore, anticoagulant therapy was required. Three patients with postoperative peroneal nerve palsy required foot orthoses. No patients complained of knee instability; however, among patients who were treated without ligamentous reconstruction, 1 patient exhibited knee varus instability when compared with the unaffected side on a varus stress test (Fig. 2).

Regarding postoperative complications, wound dehiscences, which take 3–36 weeks for full recovery, occurred in 2 of 12 patients. Local recurrence (2/12) occurred in GCT patients who were treated with curettage and adjuvant therapy. As mentioned above, 2 local recurrences of GCT were treated with marginal excision of the fibular head. Lung metastases occurred in 1 OS patient and 1 MFH patient.

4. Discussion

Tumors of the proximal fibula are rare. Because of the anatomical characteristics of this location, varus instability, peroneal

Table 1

Summary of patient demographics, diagnosis, treatment, complication, oncologic and functional outcome.

A	B	C	D	E	F	G	H	I	J	K	L	M
1	15	M	FS	13.4	CDF	100	Intentional Marginal	+	–	–	–	Suture anchor
2	17	M	OS	10.2	CDF	100	Intentional Marginal	+	–	–	–	Spike washer
3	11	M	OS	9.9	CDF	100	Intentional Marginal	+	–	–	–	Suture
4	14	F	OS	5.3	DOD	100	Intentional Marginal	+	–	–	–	Spike washer
5	49	M	CS	3.7	CDF	93	Marginal	+	–	–	–	Spike washer
6	21	M	GCT	3.7	NED	93	Marginal	+	–	–	–	Spike washer
7	30	F	GCT	5.1	NED	100	Marginal	+	+	–	–	–
8	49	F	MFH	5.4	CDF	70	Wide	–	–	40	Peroneal nerve palsy	–
9	58	F	GCT	2.8	CDF	70	Curettage	+	+	–	Peroneal nerve palsy and DVT	Intact
10	24	M	GCT	4.7	CDF	100	Curettage	+	+	–	Wound infection	Intact
11	13	M	EWS	10.1	CDF	100	Intentional Marginal	+	–	30	–	–
12	71	F	MFH	2.7	AWD	60	Wide	–	–	40	Peroneal nerve palsy and wound dehiscence	–

A – Patient

B – Age

C – Sex

D – Diagnosis. FS: Fibrosarcoma, OS: Osteosarcoma, CS: Chondrosarcoma (low-grade),

GCT: Giant cell tumor, MFH: Malignant fibrous histiocytoma, EWS: Ewing's sarcoma

E – Follow up period (years)

F – Outcome. CDF: continuous disease free, NED: no evidence of disease, AWD: alive with disease

G – MSTs Score (%)

H – Operative method

I – Attempt to preserve peroneal nerve

J – Intra-operative adjuvant treatment: A; burring, liquid nitrogen spray, alcohol

K – Postoperative radiotherapy (Gy)

L – Complication. DVT: deep venous thrombosis

M – Reconstruction of Lateral Collateral Ligament.

Download English Version:

<https://daneshyari.com/en/article/5527645>

Download Persian Version:

<https://daneshyari.com/article/5527645>

[Daneshyari.com](https://daneshyari.com)