



Mitochondrial respiratory chain disorganization in Parkinson's disease-relevant PINK1 and DJ1 mutants



Irene Lopez-Fabuel^{a, b}, Lucia Martin-Martin^{a, b}, Monica Resch-Beusher^{a, b},
Garikoitz Azkona^c, Rosario Sanchez-Pernaute^d, Juan P. Bolaños^{a, b, *}

^a University of Salamanca, CIBERFES, Institute of Functional Biology and Genomics (IBFG), CSIC, 37007 Salamanca, Spain

^b Institute of Biomedical Research of Salamanca (IBSAL), University Hospital of Salamanca, 37007 Salamanca, Spain

^c Animal Research Facility, Scientific and Technological Centers, University of Barcelona, 08007 Barcelona, Spain

^d Iniciativa Andaluza en Terapias Avanzadas, 41006 Sevilla, Spain

ARTICLE INFO

Article history:

Received 31 January 2017

Received in revised form

21 March 2017

Accepted 24 March 2017

Available online 11 April 2017

Keywords:

PINK1

DJ1

Human

Mouse

Neurons

Parkinson's disease

Mitochondria

Complexes

ABSTRACT

Brain mitochondrial complex I (CI) damage is associated with the loss of the dopaminergic neurons of the *Substantia Nigra* in Parkinson's Disease (PD) patients. However, whether CI inhibition is associated with any alteration of the mitochondrial respiratory chain (MRC) organization in PD patients is unknown. To address this issue, here we analyzed the MRC by blue native gel electrophoresis (BNGE) followed by western blotting, in mitochondria purified from fibroblasts of patients harboring PD-relevant *Pink1* mutations. We found a decrease in free CI, and in free versus supercomplexes (SCs)-assembled CI in PD; however, free complex III (CIII) was only modestly affected, whereas its free versus SCs-assembled forms decreased. Interestingly, complex IV (CIV) was considerably lost in the PD samples. These results were largely confirmed in mitochondria isolated from cultured neurons from *Pink1*^{-/-} mice, and in cultured neurons and forebrain samples from the PD-related *Dj1*^{-/-} mice. Thus, besides CI damage, the MRC undergoes a profound structural remodeling in PD likely responsible for the energetic inefficiency and mitochondrial reactive oxygen species (mROS) over-production observed in this disease.

© 2017 Elsevier Ltd. All rights reserved.

1. Introduction

The MRC complexes can be organized into SCs in a process that has been proposed to regulate electron transfer efficiency (Bianchi et al., 2004) and mROS production (Bianchi et al., 2004; Lapuente-Brun et al., 2013). Recently, we reported that the extent of assembly of CI (NADH-ubiquinone oxidoreductase) with CIII (ubiquinol-cytochrome c oxidoreductase)-containing SCs differs between neurons and astrocytes (Lopez-Fabuel et al., 2016, 2017), determining differences in mitochondrial metabolism and mROS generation

between these cells. Thus, in neurons CI is predominantly assembled into I-III SCs, whereas in astrocytes the abundance of free CI is higher (Lopez-Fabuel et al., 2016). Therefore, it can be hypothesized that changes in the normal MRC organization under certain brain pathological situations might impact on energy efficiency, mROS and neuronal survival.

It is well known that mitochondrial CI is damaged in the *Substantia Nigra* of PD patients (Schapira et al., 1989, 1990). Mutations in PTEN (phosphatase and tensin homologue)-induced kinase 1 (PINK1), a mitochondrial Ser/Thr kinase, cause an autosomal recessive form of PD (Valente et al., 2004). By phosphorylating PARKIN, PINK1 regulates the turnover of damaged mitochondria (Clark et al., 2006; Park et al., 2006). Interestingly, in PINK1 knockout (*Pink1*^{-/-}) mice it has been described that CI subunit NDUFA10 undergoes PINK1-mediated Ser²⁵⁰ phosphorylation to support CI-mediated ubiquinone reductive activity (Morais et al., 2009, 2014). Furthermore, in *Pink1*^{-/-} flies, a loss in CI assembly has been observed along with a loss in CIV (cytochrome c oxidase) activity (Liu et al., 2011). However, whether these changes alter the MRC supercomplex organization in PD patients is unknown. Here,

Abbreviations: AO, with antioxidants; BNGE, blue native gel electrophoresis; CI, complex I; CIII, complex III; CIV, complex IV; DMEM, Dulbecco's Modified Eagle's Medium; FCS, fetal calf serum; MRC, mitochondrial respiratory chain; mROS, mitochondrial reactive oxygen species; PAGE, polyacrylamide gel electrophoresis; PBS, phosphate buffered saline; Scaf1, supercomplex assembly factor 1; SC, supercomplex.

* Corresponding author. Institute of Functional Biology and Genomics (IBFG), University of Salamanca-CSIC, Zacarias Gonzalez, 2, 37007 Salamanca, Spain.

E-mail address: jbolanos@usal.es (J.P. Bolaños).

we addressed this issue and found that the MRC in fibroblasts from PD patients harboring *Pink1* mutations, and in primary neurons from *Pink1*^{-/-} mice, was disorganized. These changes were confirmed in primary neurons and brain tissue from the PD-relevant *Dj1*^{-/-} mice.

2. Materials and methods

2.1. Human samples

Skin samples were obtained from subjects expressing mutated forms of PINK1 (samples PDP1, PDP3 and PDP7) diagnosed at the Hospital Universitario Insular de Gran Canaria (La Palma de Gran Canaria, Spain), Hospital San Cecilio (Granada, Spain) and from age-matched healthy individuals (samples FH1103, FH0819 and FH0821) at the Hospital Donostia and Onkologikoa (San Sebastian, Spain) (Azkona et al., 2016). PD patients presented an early-onset, typical parkinsonian syndrome, characteristic of PINK1-associated PD (Samaranch et al., 2010). Demographic data are provided in Azkona et al. (2016). Dermal fibroblasts were cultivated as described previously (Lopez de Maturana et al., 2014) in Dulbecco's Modified Eagle's Medium (DMEM) containing 25 mM glucose and 10% (vol/vol) of fetal calf serum. Fibroblasts were used at passage 12–23.

2.2. Protocol approvals, registrations, and patient consents

The study was approved by the Ethical Committee on the Use of Human Subjects in Research in Euskadi, Spain. All subjects gave informed consent for the study using forms approved by the ethical committees on the Use of Human Subjects in Research at Hospital Universitario Insular de Gran Canaria, La Palmas de Gran Canaria; Hospital Donostia and Onkologikoa, San Sebastián; and Hospital San Cecilio, Granada, respectively.

2.2.1. Ethical statement regarding the use of animals

Mice were bred at the Animal Experimentation Unit of the University of Salamanca, and all protocols were approved by the Bioethics Committee of the University of Salamanca in accordance with the Spanish legislation (RD53/2013). PINK1 knockout (*Pink1*^{-/-}) mice (Requejo-Aguilar et al., 2014) and DJ1 knockout (*Dj1*^{-/-}) mice (Requejo-Aguilar et al., 2015) were bred under a C57Bl/6 background under heterozygosis. *Pink1*^{-/-}, *Dj1*^{-/-} and their corresponding wild type (*Pink1*^{+/+} or *Dj1*^{+/+}) siblings were used as controls.

2.2.2. Neurons in primary culture

PINK1-, DJ1-knockout, or wild type mice cortical primary neurons (Requejo-Aguilar et al., 2014, 2015) were prepared from 15.5 days-old embryos, seeded at 2.0×10^5 cells/cm² in plastic plates coated with poly-D-lysine (10 µg/ml) and incubated in Neurobasal (Life Technologies) supplemented with 2 mM of glutamine and 2% (v/v) B27-with antioxidants (AO) supplement. Cells were incubated at 37 °C in a humidified 5% (v/v) CO₂-containing atmosphere. At 72 h after plating, medium was replaced, and cells used at day 7.

2.2.3. Mitochondria isolation and solubilization

Mitochondria were obtained according to a previously published protocol (Acin-Perez et al., 2008). Briefly, cells (12–100 millions) were collected, and cell pellets frozen at –80 °C. Forebrain tissue or cell pellets (neurons or dermal fibroblasts) were homogenized (10 strokes) in a glass-teflon Potter-Elvehjem, in Buffer A (sacrose 83 mM; MOPS 10 mM; pH 7.2). The same volume of Buffer B (sacrose 250 mM; MOPS 30 mM) was added to the sample, and the homogenate was centrifuged (1 000 g, 5 min) to remove unbroken cells and nuclei. Centrifugation of the supernatant was then

performed (12,000 g, 2 min) to obtain the mitochondrial fraction, which was washed in Buffer C (sacrose 320 mM; EDTA 1 mM; Tris-HCl 10 mM; pH 7.4). Mitochondria were suspended in Buffer D (6-aminohexanoic acid 1 M; Bis-Tris-HCl 50 mM; pH 7.0). Solubilization of mitochondria was performed with digitonin at 4 g/g (5 min in ice). After a 30 min centrifugation at 13,000g, the supernatant was collected.

2.2.4. Blue native gel electrophoresis (BNGE) and western blotting

For the assessment of mitochondrial complexes organization, digitonin solubilized mitochondria (5–20 µg) were loaded in NativePAGE Novex 3–12% gels (Life Technologies). After electrophoresis, in-gel NADH dehydrogenase activity was evaluated (Diaz et al., 2009), or the resolved proteins were transferred electrophoretically to nitrocellulose membranes (Amersham protran premium 0.45 nitrocellulose, Amersham). Membranes were blocked with 5% (w/v) low-fat milk in 20 mM Tris, 150 mM NaCl, and 0.1% (w/v) Tween 20, pH 7.5, for 1 h. Subsequently of blocking, membranes were immunoblotted with NDUFS1 (a CI subunit) (1/500 by vol; sc-50132; Santa Cruz Biotechnologies), UQCRC2 (a CIII subunit) (1/1 000 by vol; ab14745; Abcam) or MT-CO1 (a CIV subunit) (1/1 000 by vol; ab14705; Abcam) primary antibody overnight at 4 °C. After incubation with horseradish peroxidase-conjugated goat anti-mouse IgG (1/10,000 by vol; 170–6 516; BioRad) or rabbit anti-goat IgG (1/10,000 by vol; ab6741; Abcam), membranes were immediately incubated with the enhanced chemiluminescence kit WesternBright ECL (Advansta), before exposure to Fuji Medical X-Ray film (Fujifilm), and the autoradiograms scanned. In all cases, at least three biologically independent replicates, or human samples, were performed, though only one representative western blot is shown in the article. The protein abundances of the protein bands were measured by densitometry on the films using ImageJ 1.48u4 software (National Institutes of Health, USA); the ratio of free versus SC-embedded CI or CIII, and free CIV versus free CIII band intensities were calculated against control samples (control or WT).

2.2.5. Protein determinations

Protein samples were quantified by the bicinchoninic acid protein assay kit (Thermo) following the manufacturer's instructions, using bovine serum albumin as a standard.

2.2.6. Statistical analysis

All measurements were carried out at least in three independent culture preparations or animals, and the results were expressed as the mean values ± SEM. The statistical analysis of the results was performed by the Student's *t*-test using the SPSS software. In all cases, *p* < 0.05 was considered significant.

3. Results

In order to investigate the MRC organization in PD, mitochondria purified from fibroblasts of patients harboring PD-relevant *Pink1* mutations, and age-matched control subjects (Azkona et al., 2016), were analyzed by BNGE followed by western blotting. We had previously shown that these cells showed increased glycolytic rate (Azkona et al., 2016), which is often a feature of damaged mitochondria. As shown in Fig. 1A, we observed a decrease in free CI, and in free versus SCs-assembled CI in the PD human fibroblasts. Free complex III (CIII) was only modestly affected, whereas the free versus SCs-assembled CIII was decreased (Fig. 1A). Interestingly, CIV protein was considerably lower in the human PD fibroblasts (Fig. 1A). To confirm these results in neural samples, we performed neurons in primary cultures from the *Pink1*^{-/-} mice, in which we also previously described increased glycolytic rate (Requejo-

Download English Version:

<https://daneshyari.com/en/article/5534673>

Download Persian Version:

<https://daneshyari.com/article/5534673>

[Daneshyari.com](https://daneshyari.com)