

Accepted Manuscript

A preliminary field trial evaluating the efficacy of 4% polyacrylamide hydrogel in horses with osteoarthritis

Scott R. McClure, Chong Wang

PII: S0737-0806(16)30596-2

DOI: [10.1016/j.jevs.2017.02.019](https://doi.org/10.1016/j.jevs.2017.02.019)

Reference: YJEVS 2286

To appear in: *Journal of Equine Veterinary Science*


Received Date: 12 October 2016

Revised Date: 8 February 2017

Accepted Date: 8 February 2017

Please cite this article as: McClure SR, Wang C, A preliminary field trial evaluating the efficacy of 4% polyacrylamide hydrogel in horses with osteoarthritis, *Journal of Equine Veterinary Science* (2017), doi: 10.1016/j.jevs.2017.02.019.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 **Short Communication**

2

3 A preliminary field trial evaluating the efficacy of 4% polyacrylamide hydrogel in horses with
4 osteoarthritis.

5

6 *Scott R. McClure^a, Chong Wang^b

7

8 ^aDepartment of Veterinary Clinical Sciences, ^bDepartments of Veterinary Diagnostic and
9 Production Animal Medicine and Department of Statistics, College of Veterinary Medicine,
10 Iowa State University, Ames, Iowa, USA.

11

12 * Corresponding author at: Midwest Equine, Boone, IA 50036.

13 *E-mail address:* scottmccluredvm@gmail.com

14

15 **Keywords**

16 Horse; Polyacrylamide; Osteoarthritis; Field Trial

17

18 **Abstract**

19 Intra-articular polyacrylamide hydrogel is being investigated for the treatment of osteoarthritis in
20 multiple species. A 4% polyacrylamide hydrogel has been evaluated in humans with positive
21 results. A field trial utilizing 4% intraarticular polyacrylamide hydrogel in equine joints with
22 naturally occurring osteoarthritis is indicated to determine the effectiveness of this therapy. The
23 objective of this field trial is to investigate the effect of intra-articular 4% polyacrylamide

Download English Version:

<https://daneshyari.com/en/article/5535630>

Download Persian Version:

<https://daneshyari.com/article/5535630>

[Daneshyari.com](https://daneshyari.com)