

Accepted Manuscript

Comparing salinities of 10, 20, and 30‰ in intensive, commercial-scale biofloc shrimp (*Litopenaeus vannamei*) production systems

Andrew J. Ray, Jeffrey M. Lotz



PII: S0044-8486(16)31191-7
DOI: doi: [10.1016/j.aquaculture.2017.03.047](https://doi.org/10.1016/j.aquaculture.2017.03.047)
Reference: AQUA 632594
To appear in: *aquaculture*
Received date: 13 December 2016
Revised date: 18 March 2017
Accepted date: 27 March 2017

Please cite this article as: Andrew J. Ray, Jeffrey M. Lotz , Comparing salinities of 10, 20, and 30‰ in intensive, commercial-scale biofloc shrimp (*Litopenaeus vannamei*) production systems. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Aqua(2017), doi: [10.1016/j.aquaculture.2017.03.047](https://doi.org/10.1016/j.aquaculture.2017.03.047)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Comparing salinities of 10, 20, and 30‰ in intensive, commercial-scale biofloc shrimp
(*Litopenaeus vannamei*) production systems

Andrew J. Ray¹, Jeffrey M. Lotz

Division of Coastal Sciences, Gulf Coast Research Laboratory, The University of Southern
Mississippi, 703 East Beach Drive, Ocean Springs, MS 39564 USA

¹Corresponding Author:

Present Address: Aquaculture Division, Kentucky State University, Land Grant Program
103 Athletic Rd., Frankfort KY, USA.

Email: andrew.ray@kysu.edu

Phone: 1-502-597-8109

Download English Version:

<https://daneshyari.com/en/article/5539419>

Download Persian Version:

<https://daneshyari.com/article/5539419>

[Daneshyari.com](https://daneshyari.com)