Accepted Manuscript



Title: Application of stable isotope analysis for detecting filial cannibalism

Authors: Atsushi Sogabe, Hideki Hamaoka, Atsushi Fukuta, Jun-ya Shibata, Jun Shoji, Koji Omori

PII: S0376-6357(16)30202-9
DOI: http://dx.doi.org/doi:10.1016/j.beproc.2017.03.019
Reference: BEPROC 3421

To appear in: *Behavioural Processes*

Received date: 18-8-2016

Please cite this article as: Sogabe, Atsushi, Hamaoka, Hideki, Fukuta, Atsushi, Shibata, Jun-ya, Shoji, Jun, Omori, Koji, Application of stable isotope analysis for detecting filial cannibalism.Behavioural Processes http://dx.doi.org/10.1016/j.beproc.2017.03.019

Application of stable isotope analysis for detecting filial cannibalism

Atsushi Sogabe[1*†], Hideki Hamaoka[2], Atsushi Fukuta[1], Jun-ya Shibata[2], Jun Shoji[1] and Koji Omori[2]

[1] Graduate School of Biosphere Science, Hiroshima University, 1-4-4 Kagamiyama, Higashi-Hiroshima 739-8528, Japan

[2] Center for Marine Environmental Studies, Ehime University, 2-5 Bunkyo-cho, Matsuyama, 790-8577, Japan

*Corresponding Author

e-mail: atsushi.sogabe@hirosaki-u.ac.jp

†Present address

Department of Biology, Faculty of Agriculture and Life Science, Hirosaki University, 3 Bunkyo-cho, Hirosaki 036-8561, Japan.

Number of words: 2820

Number of figures: 1

Number of tables: 1